

**INTERNATIONAL CONFERENCE ON TORIC TOPOLOGY  
OSAKA CITY UNIVERSITY  
29 MAY - 3 JUNE OSAKA 2006**

The homotopy type of the complement  
of a coordinate subspace arrangement

Jelena Grbić  
University of Aberdeen

Problem

**COORDINATE SUBSPACE ARRANGEMENT**

is a finite set  $\mathcal{CA} = \{L_1, \dots, L_r\} \subset \mathbb{C}^n$  of coordinate subspaces, that is,

$$L_\omega = \{(z_1, \dots, z_n) \in \mathbb{C}^n : z_{i_1} = \dots = z_{i_k} = 0\},$$

where  $\omega = \{i_1, \dots, i_k\} \subset [n]$  and its complement  $U(\mathcal{CA})$  is defined as

$$U(\mathcal{CA}) := \mathbb{C}^n \setminus \bigcup_{i=1}^r L_i.$$

**GOAL:** The homotopy type of  $U(\mathcal{CA})$ .

Toric topology-main definitions and constructions

**SIMPLICIAL COMPLEXES**

$V = \{v_1, \dots, v_n\} = [n]$  set of vertices

$K := \{\sigma_1, \dots, \sigma_s : \sigma_i \subset V\} (\emptyset \in K)$  – abstract simplicial complex closed under formation of subsets

$\sigma \in K$  – simplex  $\dim(K) = d$  if  $\#\sigma \leq d + 1$  for all  $\sigma \in K$

**STANLEY-REISNER FACE RING**

$R$  – commutative ring with unit;

$\deg(v_i) = 2$  – topological grading

$R[V] = R[v_1, \dots, v_n]$  graded polynomial algebra on  $V$  over  $R$

Given  $\sigma \subset [n]$ , set

$$v^\sigma := \prod_{i \in \sigma} v_i, \quad v^\sigma = v_{i_1} \dots v_{i_r} \quad \text{for } \sigma = \{i_1, \dots, i_r\}.$$

The **Stanley-Reisner algebra (or face ring) of  $K$**  is

$$R[K] := R[v_1, \dots, v_n] / (v^\sigma : \sigma \notin K).$$

''Topological models'' for the algebraic objects

### DAVIS–JANUSZKIEWICZ SPACE $DJ(K)$

–topological realisation of the Stanley–Reisner ring  $R[K]$ , that is,

$$H^*(DJ(K); R) = R[K] \quad (\text{for } R = \mathbb{Z} \text{ or } R = \mathbb{Z}/2).$$

Davis–Januszkiewicz  $DJ(K) = ET^n \times_{T^n} \mathcal{Z}_K$

Buchstaber–Panov through a simple colimit of nice blocks

Assume  $R = \mathbb{Z}$ . Denote  $\mathbb{C}P^\infty = BS^1$ , thus  $BT^n = (\mathbb{C}P^\infty)^n$

For  $\omega \subset [n]$ , define

$$BT^\omega := \{(x_1, \dots, x_n) \in BT^n : x_i = * \text{ if } i \notin \omega\}.$$

For  $K$  on  $[n]$ , the Davis-Januszkiewicz space of  $K$  is given by

$$DJ(K) := \bigcup_{\sigma \in K} BT^\sigma \subset BT^n.$$

### MOMENT–ANGLE COMPLEX $\mathcal{Z}_K$

Torus  $T^n \subset (D^2)^n = \{(z_1, \dots, z_n) \in \mathbb{C}^n : |z_i| \leq 1, \forall i\}$

For arbitrary  $\sigma \subset [n]$ , define

$$B_\sigma := \{(z_1, \dots, z_n) \in (D^2)^n : |z_i| = 1 \text{ if } i \notin \sigma\}.$$

$$B_\sigma \cong (D^2)^{|\sigma|} \times T^{n-|\sigma|}$$

For  $K$  on  $[n]$ , define the moment–angle complex  $\mathcal{Z}_K$  by

$$\mathcal{Z}_K := \bigcup_{\sigma \in K} B_\sigma \subset (D^2)^n.$$

$B_\sigma$  invariant under the action of  $T^n \rightsquigarrow T^n$  acts on  $\mathcal{Z}_K$

**Proposition.** *The moment–angle complex  $\mathcal{Z}_K$  is the homotopy fibre of the inclusion*

$$i: \text{DJ}(K) \longrightarrow BT^n.$$

**Proposition.**  $H_{T^n}^*(\mathcal{Z}_K) = \mathbb{Z}[K]$

Arrangements and their complements

For  $K$  on set  $[n]$ , define the complex coordinate subspace arrangement as

$$\mathcal{CA}(K) := \{L_\sigma : \sigma \notin K\}$$

and its complement in  $\mathbb{C}^n$  by

$$U(K) := \mathbb{C}^n \setminus \bigcup_{\sigma \notin K} L_\sigma.$$

If  $L \subset K$  is a subcomplex, then  $U(L) \subset U(K)$ .

**Proposition.** *The assignment*

$$K \mapsto U(K)$$

*defines a one–to–one order preserving correspondence*

$$\left\{ \begin{array}{c} \text{simplicial} \\ \text{complexes on } [n] \end{array} \right\} \longleftrightarrow \left\{ \begin{array}{c} \text{complements of} \\ \text{coordinate subspace} \\ \text{arrangements in } \mathbb{C}^n \end{array} \right\}.$$

## CONNECTION BETWEEN $\mathcal{Z}_K$ AND $U(K)$

**Theorem** (Buchstaber–Panov). *There is an equivariant deformation retraction*

$$U(K) \xrightarrow{\cong} \mathcal{Z}_K.$$

## COHOMOLOGY OF $U(K)$

**Theorem** (Buchstaber–Panov). *The following isomorphism of graded algebra holds*

$$H^*(U(K); k) \cong \mathrm{Tor}^{k[v_1, \dots, v_n]}(k[K], k) \cong H[\wedge[u_1, \dots, u_n] \otimes k[K], d].$$

hints from ALGEBRA and COMBINATORICS

**Definition.** The Stanley-Reisner ring  $k[K]$  is **Golod** if all Massey products in  $\mathrm{Tor}^{k[v_1, \dots, v_n]}(k[K], k)$  vanish.

**Definition.** A simplicial complex  $K$  is **shifted** if there is an ordering  $\sigma \in K$ ,  $v' < v \Rightarrow (\sigma - v) \cup v' \in K$ .

**Proposition.** *If  $K$  is shifted, then its face ring  $k[K]$  is Golod.*

## THE MAIN THEOREM (G., Theriault)

Let  $K$  be a shifted complex. Then  $\mathcal{Z}_K$  is a wedge of spheres.

Back to COMBINATORICS

**PROBLEM:** Determine the homotopy type of the complement of arbitrary codimension coordinate subspace arrangements.

**STRATEGY:**

- 1) determine the simplicial complex  $K$  which corresponds to a codimension- $i$  coordinate subspace arrangement,  $U(K)$ ;
- 2) associate to the determined simplicial complex  $K$  its Davis-Januszkiewicz space, i.e,  $DJ(K)$ ;
- 3) looking at the fibration

$$\mathcal{Z}_K \longrightarrow DJ(K) \longrightarrow BT^n,$$

describe the homotopy type of  $\mathcal{Z}_K$ .

**1)** Look at an  $i+2$ -codimension coordinate subspace in  $\mathbb{C}^n$ , that is,  $L_\omega = \{(z_1, \dots, z_n) \in \mathbb{C}^n : z_{j_1} = \dots = z_{j_{i+2}} = 0\}$ ,  $\omega = \{j_1, \dots, j_{i+2}\}$ .

Then  $K = \text{sk}^i(\Delta^{n-1})$ .

Hence,  $\mathbb{C}^n \setminus \mathcal{CA}^{i+2} = U(\text{sk}^i(\Delta^{n-1}))$ .

**2)** A colimit model of the Davis-Januszkiewicz space for  $K$  is given by

$$DJ(K) := \bigcup_{\sigma \in K} BT^\sigma \subset BT^n, \quad \# \text{vertices in } K.$$

Then we have

$$\begin{aligned} DJ(K) &= T_{n-1-i}^n \\ &= \{(z_1, \dots, z_n) : \text{at least } n-1-i \text{ coordinates are } *\} \subset (\mathbb{C}P^\infty)^n. \end{aligned}$$

3) Determine the homotopy fibre  $\mathcal{Z}_K$  of the fibration sequence

$$(\mathcal{Z}_K)_k^n \longrightarrow T_k^n \longrightarrow (\mathbb{C}P^\infty)^n \text{ for } 1 \leq k \leq n - 1.$$

Let  $X_1, \dots, X_n$  be path-connected spaces.

There is a filtration of  $X_1 \times \dots \times X_n$  given by

$$T_n^n \longrightarrow T_{n-1}^n \longrightarrow \dots \longrightarrow T_0^n$$

where  $T_k^n = \{(x_1, \dots, x_n) \in X_1 \times \dots \times X_n : \text{at least } k \text{ of } x_i\text{'s are } *\}$ .

**Theorem** (Porter; G., Theriault). *For  $n \geq 1$ , and  $k$  such that  $1 \leq k \leq n - 1$ , the homotopy fibre  $F_k^n$  of the inclusion  $i: T_k^n \longrightarrow X_1 \times \dots \times X_n$  decomposes as*

$$F_k^n \simeq \bigvee_{j=n-k+1}^n \left( \bigvee_{1 \leq i_1 < \dots < i_j \leq n} \binom{j-1}{n-k} \Sigma^{n-k} \Omega X_{i_1} \wedge \dots \wedge \Omega X_{i_j} \right).$$

□

Take for  $X_1 = \dots = X_n = \mathbb{C}P^\infty$ . Then we have the inclusion

$$i: T_k^n(\mathbb{C}P^\infty) \longrightarrow (\mathbb{C}P^\infty)^n.$$

It follows that

$$\begin{aligned} F_k^n &\simeq \bigvee_{j=n-k+1}^n \left( \binom{n}{j} \binom{j-1}{n-k} \Sigma^{n-k} \underbrace{\Omega \mathbb{C}P^\infty \wedge \dots \wedge \Omega \mathbb{C}P^\infty}_j \right) \\ &\simeq \bigvee_{j=n-k+1}^n \binom{n}{j} \binom{j-1}{n-k} S^{n+j-k}. \end{aligned}$$

## Family

$$\mathcal{F}_t = \{K - \text{simplicial complex} \mid \Sigma^t \mathcal{Z}_K \text{ a wedge of spheres}\}$$

Notice that  $\mathcal{F}_0 \subset \mathcal{F}_1 \subset \dots \subset \mathcal{F}_t \subset \dots \subset \mathcal{F}_\infty$ .

We have shown if  $K$  -shifted, then  $K \in \mathcal{F}_0$  ( $sk^i(\Delta^{n-1}) \in \mathcal{F}_0$ )

Want: make simplicial complexes out of our building blocks

### WHAT CAN HOMOTOPY SEE?

#### DISJOINT UNION OF SIMPLICIAL COMPLEXES

Let  $K_1 \in \mathcal{F}_t$  and  $K_2 \in \mathcal{F}_s$ . Then  $K_1 \amalg K_2 \in \mathcal{F}_m$ ,  $m = \max\{t, s\}$ .

$$\mathcal{Z}_K \simeq (\prod_i S^1 * \prod_j S^1) \vee (\mathcal{Z}_{K_1} \times \prod_i S^1) \vee (\prod_j S^1 \times \mathcal{Z}_{K_2})$$

#### GLUING ALONG A COMMON FACE

Let  $K = K_1 \cup_\sigma K_2$ . If  $K_1, K_2 \in \mathcal{F}_0$ , then  $K \in \mathcal{F}_0$ .

$$\mathcal{Z}_K \simeq (\prod S^1 * \prod S^1) \vee (\mathcal{Z}_{K_1} \times \prod S^1) \vee (\mathcal{Z}_{K_2} \times \prod S^1)$$

#### JOIN OF SIMPLICIAL COMPLEXES

$K_1, K_2$  simplicial complexes on sets  $S_1$  and  $S_2$ , belonging to  $\mathcal{F}_t$  and  $\mathcal{F}_s$ .

The *join*  $K_1 * K_2 := \{\sigma \subset S_1 \cup S_2 : \sigma = \sigma_1 \cup \sigma_2, \sigma_1 \in K_1, \sigma_2 \in K_1\}$

Notice  $k[K_1 * K_2] = k[K_1] \otimes k[K_2]$

Therefore for the join of  $K_1$  and  $K_2$  we get a product fibration

$$DJ(K_1) \times DJ(K_2) \longrightarrow BT^{m_1} \times BT^{m_2}$$

hence  $\mathcal{Z}_{K_1 * K_2} \simeq \mathcal{Z}_{K_1} \times \mathcal{Z}_{K_2}$  and  $K_1 * K_2 \in \mathcal{F}_m$ ,  $m = \max\{t, s\} + 1$ .

Our contribution to ALGEBRA

Let  $A$  be a polynomial ring on  $n$  variables  $k[x_1, \dots, x_n]$  over a field  $k$  and  $S = A/I$ , where  $I$  is a homogeneous ideal, i.e,  $S = k[K]$  for some simplicial complex  $K$ .

**PROBLEM:** The nature of  $\text{Tor}^S(k, k)$ .

The Poincaré series

$$P(S) = \sum_{i=0}^{\infty} b_i t^i \quad \text{where } b_i = \dim_k \text{Tor}_i^S(k, k)$$

**PROBLEM:** The rationality of  $P(S)$ .

**Theorem.** (Golod) *There exist non-negative integers  $n, c_1, \dots, c_n$  such that*

$$P(S) \leq \frac{(1+t)^n}{1 - \sum_{i=1}^n c_i t^{i+1}}.$$

**Theorem.** (G., Theriault) *There is a topological proof of Golod's inequality.*

**Theorem.** (*Buchstaber-Panov-Ray*)

$$\mathrm{Tor}_*^{k[K]}(k, k) \cong H^*(\Omega DJ(K); k).$$

Looking at the split fibration

$$\Omega \mathcal{Z}_K \longrightarrow \Omega DJ(K) \longrightarrow T^n$$

$$\mathrm{Tor}_*^{k[K]}(k, k) \cong H^*(\Omega DJ(K)) = H^*(T^n) \otimes H^*(\Omega \mathcal{Z}_K)$$

Using the bar resolution,

$$P(H^*(\Omega \mathcal{Z}_K)) \leq P(T(\Sigma^{-1} H^*(\mathcal{Z}_K))).$$

Therefore

$$\begin{aligned} P(k[K]) &= (1+t)^n P(H^*(\Omega \mathcal{Z}_K)) \leq (1+t)^n P(T(\Sigma^{-1} H^*(\mathcal{Z}_K))) \\ &= \frac{(1+t)^n}{1 - P(\Sigma^{-1} H^*(\mathcal{Z}_K))}. \end{aligned}$$

Equality is obtained when  $H^*(\mathcal{Z}_K)$  is Golod.

**Corollary.** *When  $K \in \mathcal{F}_0$ , then  $P(k[K])$  is rational.*