MOMENT-ANGLE COMPLEXES AND COMBINATORICS OF SIMPLICIAL MANIFOLDS

VICTOR M. BUCHSTABER AND TARAS E. PANOV

Let $\rho : (D^2)^m \to I^m$ be the orbit map for the diagonal action of torus T^m on the unit poly-disk $(D^2)^m \subset \mathbb{C}^m$. Each face of the cube $I^m = [0,1]^m$ (viewed as a cubical complex) has the form

$$F_{I\subset J} = \{(y_1, \dots, y_m) \in I^m \ : \ y_i = 0 \text{ if } i \in I, \ y_j = 1 \text{ if } j \notin J\},$$

where $I \subset J$ are two subsets of the index set $[m] = \{1, \dots, m\}$. For each face $F_{I\subset J}$ put $B_{I\subset J} := \rho^{-1}(F_{I\subset J})$. If $\#I = i$, $\#J = j$, then $B_{I\subset J} \cong (D^2)^{j-i} \times T^{m-j}$.

Definition 1. Let C be a cubical subcomplex of I^m. The *moment-angle complex* $\mathrm{ma}(C)$ is the T^m-invariant decomposition of the subset $\rho^{-1}(|C|) \subset (D^2)^m$ into blocks $B_{I\subset J}$ corresponding to the faces $F_{I\subset J}$ of complex C.

Many combinatorial problems concerning cubical complexes may be treated by studying the equivariant topology of moment-angle complexes. In the present paper we realize this approach in the case of cubical complexes determined by simplicial complexes. Let K^{n-1} be an $(n-1)$-dimensional simplicial complex with m vertices, and $|K|$ the corresponding polyhedron. If $I = \{i_1, \dots, i_k\} \subset [m]$ is a simplex of K, then we would write $I \in K$. Define the following two cubical subcomplexes of I^m:

$$\mathrm{cub}(K) = \{F_{I\subset J} \ : \ J \in K, I \neq \varnothing\}, \quad \mathrm{cc}(K) = \{F_{I\subset J} \ : \ J \in K\}.$$

Lemma 2. *As a topological space, the complex* $\mathrm{cub}(K)$ *is homeomorphic to* $|K|$*, while* $\mathrm{cc}(K)$ *is homeomorphic to the cone* $|\mathrm{cone}(K)|$.

The cubical complex $\mathrm{cc}(K)$ was introduced in [1] and then studied in [2]. The cubical complex $\mathrm{cub}(K)$ appeared in [3].



Denote the moment-angle complexes corresponding to $\mathrm{cub}(K)$ and $\mathrm{cc}(K)$ by $\mathcal{W}_K$ and $\mathcal{Z}_K$ respectively. Consider the cellular decomposition of the poly-disk $(D^2)^m$ that is obtained by subdividing each factor D^2 into 0-dimensional cell 1, 1-dimensional cell T, and 2-dimensional cell D, see Fig. a). Each cell of $(D^2)^m$ is a product of cells D_i, T_i, 1_i, $i = 1, \dots, m$, i.e., can be written as $D_I T_J 1_{[m]\setminus I\cup J}$, where I, J are disjoint subsets of $[m]$. Set $D_I T_J := D_I T_J 1_{[m]\setminus I\cup J}$. Now it can be easily seen that $\mathcal{Z}_K$ is cellular subcomplex of $(D^2)^m$ consisting of all cells $D_I T_J$ such that $I \in K$.

Lemma 3. *The embedding* $T^m = \rho^{-1}(1, \dots, 1) \hookrightarrow \mathcal{Z}_K$ *is a cellular map homotopic to the map to a point.*

As it was shown in [2], for any field $\mathbf{k}$ there is the following isomorphism of algebras:

$$H^*(\mathcal{Z}_K) \cong \mathrm{Tor}_{[v_1,\dots,v_m]}\big(\mathbf{k}(K), \mathbf{k}\big) = H^*\big[\mathbf{k}(K) \otimes \Lambda[u_1, \dots, u_m], d\big], \tag{1}$$

Partially supported by the Russian Foundation for Fundamental Research grant no. 99-01-00090.

where $\mathbf{k}(K)$ is the Stanley–Raisner ring of complex K, and the differential d is defined by $d(v_i)=0$, $d(u_i)=v_i$, $i=1,\dots,m$. The Tor-algebra from (1) is naturally a *bigraded* algebra with $\operatorname{bideg}(v_i)=(0,2)$, $\operatorname{bideg}(u_i)=(-1,2)$. The calculation of the ring $H^*(\mathcal{Z}_K)$ allowed to describe the multiplicative structure in the cohomology of the complement of a coordinate subspace arrangement in $\mathbb{C}^m$ [2].

In [1] there was introduced the subcomplex $\mathcal{C}^*(K)\subset \mathbf{k}(K)\otimes\Lambda[u_1,\dots,u_m]$ spanned by monomials u_J and v_Iu_J such that $I\cap J=\varnothing$, $I\in K$. It was also shown there that the cohomology of $\mathcal{C}^*(K)$ is also isomorphic to that of $\mathcal{Z}_K$. Denote by $\mathcal{C}_*(\mathcal{Z}_K)$ and $\mathcal{C}^*(\mathcal{Z}_K)$ the chain and the cochain complex for type a) cellular decomposition of $\mathcal{Z}_K$ respectively.

Theorem 4. *Let $(D_IT_J)^*\in\mathcal{C}^*(\mathcal{Z}_K)$ denote the cellular cochain dual to the cell $D_IT_J\in\mathcal{Z}_K$. The correspondence $v_Iu_J\mapsto(D_IT_J)^*$ defines a canonical isomorphism of complexes $\mathcal{C}^*(K)$ and $\mathcal{C}^*(\mathcal{Z}_K)$, each of which calculates $H^*(\mathcal{Z}_K)$.*

The pair $(\mathcal{Z}_K,T^m)$ acquires a bigraded cellular structure by setting $\operatorname{bideg}(D_i)=(0,2)$, $\operatorname{bideg}(T_i)=(-1,2)$, $\operatorname{bideg}(1_i)=(0,0)$. Put $b_{-q,2p}(\mathcal{Z}_K,T^m)=\dim H_{-q,2p}[\mathcal{C}_*(\mathcal{Z}_K,T^m)]$. Consider the new cellular decomposition of the poly-disc $(D^2)^m$ that is obtained by subdividing each factor D^2 into 5 cells D, T, I, 1, 0, see Fig. b). This allows to introduce a bigraded cellular structure on $\mathcal{W}_K$ and define the numbers $b_{q,2p}(\mathcal{W}_K)=\dim H_{q,2p}[\mathcal{C}_*(\mathcal{W}_K)]$. Put

$$\chi(\mathcal{Z}_K,T^m;t)=\sum_{p,q}(-1)^q b_{-q,2p}(\mathcal{Z}_K,T^m)t^{2p},\quad \chi(\mathcal{W}_K;t)=\sum_{p,q}(-1)^q b_{q,2p}(\mathcal{W}_K)t^{2p}.$$

Let f_i be the number of i-simplices of K, and $(h_0,\dots,h_n)$ the *h-vector* of K determined from the equation $h_0t^n+\ldots+h_{n-1}t+h_n=(t-1)^n+f_0(t-1)^{n-1}+\ldots+f_{n-1}$.

Theorem 5. *Put $h(t)=h_0+h_1t+\cdots+h_nt^n$. Then*

$$\chi(\mathcal{Z}_K,T^m;t)=(1-t^2)^{m-n}h(t^2)-(1-t^2)^m,$$
$$\chi(\mathcal{W}_K;t)=(1-t^2)^{m-n}h(t^2)+(-1)^{n-1}h_n(1-t^2)^m.$$

Lemma 6. *If $|K|\cong S^{n-1}$ (i.e., K is a simplicial sphere), then $\mathcal{Z}_K$ is a closed manifold.*

Suppose now that K^{n-1} is a simplicial manifold. Then the complex $\mathcal{Z}_K$ generally fails to be a manifold. However, removing from $\mathcal{Z}_K$ a small neighbourhood $U_\varepsilon(T^m)$ of the orbit $\rho^{-1}(1,\dots,1)\cong T^m$ we obtain manifold $W_K=\mathcal{Z}_K\setminus U_\varepsilon(T^m)$ with boundary $\partial W_K=|K|\times T^m$.

Theorem 7. *The manifold (with boundary) W_K is equivariantly homotopy equivalent to the complex $\mathcal{W}_K$. There is a canonical homeomorphism of pairs $(W_K,\partial W_K)\to(\mathcal{Z}_K,T^m)$.*

The relative Poincaré duality isomorphisms for W_K imply

$$\chi(\mathcal{W}_K;t)=(-1)^{m-n}t^{2m}\chi(\mathcal{Z}_K,T^m;\tfrac{1}{t}).$$

Taking into account Theorem 5 we obtain

Corollary 8. *Let K^{n-1} be a simplicial manifold. Then*

$$h_{n-i}-h_i=(-1)^i(h_n-1)\tbinom{n}{i}=(-1)^i\big(\chi(K^{n-1})-\chi(S^{n-1})\big)\tbinom{n}{i},\quad i=0,1,\dots,n,$$

where $\chi(\cdot)$ denotes the Euler number.

Rewriting the equation (8) in terms of the f-vector we come to more complicated equations, which were deduced in [4], [5]. For $|K|=S^{n-1}$ Corollary 8 gives the classical Dehn–Sommerville equations. In the particular case of PL-manifolds the topological invariance of numbers $h_{n-i}-h_i$ (which follows directly from Corollary 8) was firstly observed by Pachner in [6, (7.11)].

The extended version of this article is `http://xxx.lanl.gov/abs/math.AT/0005199`.
The authors wish to express special thanks to Oleg Musin for stimulating discussions and helpful comments, in particular, for drawing our attention to the results of [4] and [6].

References

[1] V. M. Buchstaber and T. E. Panov. Proceedings of the Steklov Institute of Mathematics **225** (1999), 87–120.
[2] V. M. Buchstaber and T. E. Panov. (Russian), Zap. Nauchn. Semin. POMI **266** (2000), 29–50.
[3] M. A. Shtan'ko and M. I. Shtogrin. Russian Math. Surveys **47** (1992), no. 1, 267–268.
[4] B. Chen, M. Yan. Proceedings of the Steklov Institute of Mathematics **221** (1998), 305–319.
[5] V. Klee. Canadian J. Math. **16** (1964), 517–531.
[6] U. Pachner. European J. Combinatorics **12** (1991), 129–145.

Department of Mathematics and Mechanics, Moscow State University, 119899 Moscow, RUSSIA
E-mail address: buchstab@mech.math.msu.su tpanov@mech.math.msu.su