

From real quadrics to polytopes via manifolds

Taras Panov

Moscow State University

CAT'09, 6–11 July 2009, Warszawa

1. From polytopes to quadrics.

\mathbb{R}^n : Euclidean vector space. Consider a convex polyhedron

$$P = \{\mathbf{x} \in \mathbb{R}^n : (\mathbf{a}_i, \mathbf{x}) + b_i \geq 0 \text{ for } 1 \leq i \leq m\}, \quad \mathbf{a}_i \in \mathbb{R}^n, b_i \in \mathbb{R}.$$

Assume $\dim P = n$, no redundant inequalities, P is bounded, and bounding hyperplanes $H_i = \{(\mathbf{a}_i, \mathbf{x}) + b_i = 0\}$, $1 \leq i \leq m$, intersect in general position at every vertex.

Then P is an n -dim **convex simple polytope** with m **facets**

$$F_i = \{\mathbf{x} \in P : (\mathbf{a}_i, \mathbf{x}) + b_i = 0\} = P \cap H_i$$

and normal vectors \mathbf{a}_i , for $1 \leq i \leq m$. At every vertex meets an n -tuple of facets.

Two polytopes are said to be **combinatorially equivalent** if their face posets are isomorphic.

We may specify P by a matrix inequality

$$P = \{\mathbf{x} : A_P \mathbf{x} + \mathbf{b}_P \geq 0\},$$

where $A_P = (a_{ij})$ is the $m \times n$ matrix of row vectors \mathbf{a}_i , and \mathbf{b}_P is the column vector of scalars b_i .

The affine injection

$$i_P : \mathbb{R}^n \longrightarrow \mathbb{R}^m, \quad \mathbf{x} \mapsto A_P \mathbf{x} + \mathbf{b}_P$$

embeds P into $\mathbb{R}_{\geq}^m = \{\mathbf{y} \in \mathbb{R}^m : y_i \geq 0\}$.

Now define the space \mathcal{Z}_P by a pullback diagram

$$\begin{array}{ccc} \mathcal{Z}_P & \xrightarrow{i_Z} & \mathbb{C}^m \\ \downarrow & & \downarrow \\ P & \xrightarrow{i_P} & \mathbb{R}^m \end{array} \quad \begin{array}{c} (z_1, \dots, z_m) \\ \downarrow \\ (|z_1|^2, \dots, |z_m|^2) \end{array}$$

Here i_Z is a T^m -equivariant embedding.

Prop 1. \mathcal{Z}_P is a smooth T^m -manifold with canonically trivialised normal bundle of $i_Z: \mathcal{Z}_P \rightarrow \mathbb{C}^m$.

Idea of proof.

- 1) Write the image $i_P(\mathbb{R}^n) \subset \mathbb{R}^m$ as the set of common solutions of $m - n$ linear equations $\sum_{k=1}^m c_{jk}(y_k - b_k) = 0$, $1 \leq j \leq m - n$;
- 2) replace every y_k by $|z_k|^2$ to get a representation of \mathcal{Z}_P as an intersection of $m - n$ real quadratic hypersurfaces:

$$\sum_{k=1}^m c_{jk} (|z_k|^2 - b_k) = 0, \quad \text{for } 1 \leq j \leq m - n.$$

- 3) check that 2) is a non-degenerate intersection, i.e. the gradient vectors are linearly independent at each point of \mathcal{Z}_P . □

\mathcal{Z}_P is called the **moment-angle manifold** corresponding to P .

In fact, the topological type of \mathcal{Z}_P depends only on the combinatorial type of P (the original construction of [Davis–Januszkiewicz]).

Write the system $\sum_{k=1}^m c_{jk}(|z_k|^2 - b_k) = 0$, $1 \leq j \leq m - n$, as

$$\mathcal{Z}_P = \{\mathbf{z} \in \mathbb{C}^m : C|\mathbf{z}|^2 = C\mathbf{b}_P\}$$

with $C = (c_{jk})$ $(m - n) \times m$ -matrix, $|\mathbf{z}|^2$ column of $|z_k|^2$.

Rows of C constitute a basis in the space of linear relations between the \mathbf{a}_i 's. That is, $CA_P = 0$ and $\text{rank } C = m - n$ (note $\text{rank } A_P = n$).

Given P , how to choose C ?

1st method:

Assume the first n facets F_1, \dots, F_n meet at a vertex, and take their normals as the basis for \mathbb{R}^n (after linear transformation). Then

$$A_P = \begin{pmatrix} E \\ A_P^* \end{pmatrix}$$

with E unit $n \times n$ -matrix and A_P^* an $(m - n) \times n$ -matrix. Then we may take

$$C = \begin{pmatrix} -A_P^* & E \end{pmatrix}.$$

(Remember $CA_P = 0$!)

This is convenient for applications in cobordism (finding quasitoric representatives in complex cobordism classes).

2nd method:

Have $\alpha_1 \mathbf{a}_1 + \dots + \alpha_m \mathbf{a}_m = 1$ with $\alpha_i > 0$.

In fact, $\alpha_i = \text{Vol } F_i$ if $|\mathbf{a}_i| = 1$.

By scaling the \mathbf{a}_i 's can always achieve $\mathbf{a}_1 + \dots + \mathbf{a}_m = 1$ and so take

$$C = \begin{pmatrix} C_1 \\ 1 \dots 1 \end{pmatrix} \quad \text{where } C_1 \text{ is } (m - n - 1) \times m.$$

By moving the origin 0 into $\text{Int } P$, get $b_i > 0$. By scaling P get $b_1 + \dots + b_m = 1$, so the last quadratic equation defining \mathcal{Z}_P is

$$|z_1|^2 + \dots + |z_m|^2 = 1.$$

Subtracting this from the first $m - n - 1$ equations, finally get

$$\mathcal{Z}_P = \begin{cases} \mathbf{z} \in \mathbb{C}^m: & C_* |\mathbf{z}|^2 = 0, \\ & |z_1|^2 + \dots + |z_m|^2 = 1 \end{cases}$$

where C_* is $(m - n - 1) \times m$.

Ex 1.

$$P = \left\{ \mathbf{x} \in \mathbb{R}^n : \begin{array}{l} x_i \geq 0, \quad i = 1, \dots, n, \\ -x_1 - \dots - x_n + 1 \geq 0 \end{array} \right\} : \quad n\text{-simplex.}$$

So $m = n + 1$, $\mathbf{a}_i = \mathbf{e}_i$ for $i = 1, \dots, n$, $\mathbf{a}_{n+1} = -\mathbf{e}_1 - \dots - \mathbf{e}_n$.

$$A_P = \begin{pmatrix} E \\ -1 \dots -1 \end{pmatrix}, \quad C = (1 \dots 1),$$

$$\mathcal{Z}_P = \{\mathbf{z} : |z_1|^2 + \dots + |z_m|^2 = 1\} = S^{2n+1},$$

and $C_* = \emptyset$.

2. From quadrics to polytopes.

After [Bosio–Meersseman].

$C_* = (\mathbf{c}_1, \dots, \mathbf{c}_m)$: $p \times m$ matrix, $0 \leq p < m$ (later set $p = m - n - 1$).

Set

$$\mathcal{L}_C = \left\{ \mathbf{z} \in \mathbb{C}^m : \begin{array}{l} \mathbf{c}_1|z_1|^2 + \dots + \mathbf{c}_m|z_m|^2 = 0 \\ |z_1|^2 + \dots + |z_m|^2 = 1 \end{array} \right\} : \text{“link of } p \text{ special real quadrics in } \mathbb{C}^m\text{”}.$$

C_* is **admissible** if \mathcal{L}_C is nonempty and nondegenerate.
(So \mathcal{L}_C is a $(2m - p - 1)$ -dimensional manifold.)

$\text{Conv}(C_*) :=$ convex hull of $\mathbf{c}_1, \dots, \mathbf{c}_m$ in \mathbb{R}^p .

Lemma 1. C_* is admissible iff

- 1) $0 \in \text{Conv}(C_*)$, and
- 2) $0 \in \text{Conv}(\mathbf{c}_i, i \in I)$ implies $|I| > p$.

Ex 2. $p = 1$, so $\mathbf{c}_i \in \mathbb{R}$. Then (2) implies $\mathbf{c}_i \neq 0$.

Assume k of \mathbf{c}_i are positive and $l = m - k$ are negative.

Then (1) implies $k > 0$, $l > 0$.

Get like $|z_1|^2 + \dots + |z_k|^2 - |z_{k+1}|^2 - \dots - |z_m|^2 = 0$:

cone over $S^{2k-1} \times S^{2l-1}$,

and $|z_1|^2 + \dots + |z_m|^2 = 1$, so $\mathcal{L}_C = S^{2k-1} \times S^{2l-1}$.

Ex 3. $p = 2$, so $\mathbf{c}_i \in \mathbb{R}^2$. Then by (1), $0 \in \text{Conv}(\mathbf{c}_1, \dots, \mathbf{c}_m)$.

(2) says that no segment joining \mathbf{c}_i contains 0.

Lemma 2. *Can always achieve odd number of points on a circle with positive weights assigned.*

Set $k(C) =$ number of i such that $\mathcal{L}_C \cap \{z_i = 0\} = \emptyset$.

Lemma 3. $\mathcal{L}_C \cong \mathcal{L}_{C'} \times T^{k(C)}$, where $\mathcal{L}_{C'} \subset \mathbb{C}^{m-k(C)}$ intersects every coordinate hyperplane.

How to get a polytope out of \mathcal{L}_C ?

\mathcal{L}_C/T^m is given by the nonnegative solutions of

$$C_*\mathbf{y} = 0, \quad y_1 + \dots + y_m = 1.$$

By nondegeneracy condition (2), this system has maximal rank. So may write its general solution as

$$y_i = (\mathbf{a}_i, \mathbf{x}) + b_i, \quad \mathbf{x} \in \mathbb{R}^{m-p-1}, \text{ with } b_i > 0.$$

Therefore,

$$\mathcal{L}_C/T^m = \{\mathbf{x} \in \mathbb{R}^{m-p-1} : (\mathbf{a}_i, \mathbf{x}) + b_i \geq 0\}.$$

Lemma 4. *This is a simple $(m - p - 1)$ -dimensional polytope P with $m - k(C_*)$ facets.*

So every $\{z_i = 0\} \cap \mathcal{L}_C = \emptyset$ gives a redundant inequality.

$P^* = \text{Conv}\left(\frac{\mathbf{a}_1}{b_1}, \dots, \frac{\mathbf{a}_m}{b_m}\right)$: polar (or dual) simplicial polytope.

Denote $\mathbf{a}'_i = \frac{\mathbf{a}_i}{b_i}$.

Lemma 5. $0 \in \text{Int Conv}(\mathbf{c}_i, i \in I)$

$\iff \text{Conv}(\mathbf{a}'_i, i \in [m] \setminus I)$ is a proper face of P^* .

In other words, $(\mathbf{c}_1, \dots, \mathbf{c}_m)$ is the Gale diagram of $(\mathbf{a}'_1, \dots, \mathbf{a}'_m)$.

Finally, we get

Thm 1. Every \mathcal{L}_C with $k(C) = 0$ is \mathcal{Z}_P for some P .

3. Topology of \mathcal{Z}_P .

F_1, \dots, F_m : facets of P .

Given $I \subset [m]$, set $P_I = \bigcup_{i \in I} F_i \subset P$.

Thm 2. $H^k(\mathcal{Z}_P) = \bigoplus_{I \subset [m]} \widetilde{H}^{k-|I|-1}(P_I)$.

Ex 4. P a 5-gon. Then $\dim \mathcal{Z}_P = 7$, and the Betti vector is

$$(1, 0, 0, 5, 5, 0, 0, 1).$$

In fact, $\mathcal{Z}_P = (S^3 \times S^4) \#^5$.

Proof of Thm 2. \mathcal{Z}_P generalises to \mathcal{Z}_K , the **moment-angle complex** defined for an arbitrary simplicial complex K on m vertices.

Given P as above, set

$$K_P = \left\{ \sigma = \{i_1, \dots, i_k\} : F_{i_1} \cap \dots \cap F_{i_k} \neq \emptyset \text{ in } P \right\},$$

the **boundary complex** of P^* . It is a sphere triangulation: $|K_P| \cong S^{n-1}$.

Then $\mathcal{Z}_{K_P} = \mathcal{Z}_P$.

By [Buchstaber-P], there is an isomorphism of (bi)graded algebras

$$\begin{aligned} H^*(\mathcal{Z}_K) &\cong \operatorname{Tor}_{\mathbb{Z}[v_1, \dots, v_m]}^{*,*}(\mathbb{Z}[K], \mathbb{Z}) \\ &\cong H\left[\Lambda[u_1, \dots, u_m] \otimes \mathbb{Z}[K]; d\right], \end{aligned}$$

where $\mathbb{Z}[K]$ is the **face ring** (or the **Stanley–Reisner ring**) of K ,
 $du_i = v_i$, $dv_i = 0$ for $1 \leq i \leq m$.

From this description follows Hochster's calculation of Tor modules in terms of **full subcomplexes** of K :

$$\mathrm{Tor}_{\mathbb{Z}[v_1, \dots, v_m]}^{-i, 2j}(\mathbb{Z}[K], \mathbb{Z}) \cong \bigoplus_{|J|=j} \widetilde{H}^{j-i-1}(K_J),$$

where K_J is the restriction of K to the subset $J \subset [m]$.

This dualises to the required description of the cohomology in the case $K = K_P$ (because P_J retracts onto K_J). \square

4. Quasitoric manifolds and cobordism.

Assume given P as above, and an integral $n \times m$ matrix

$$\Lambda = \begin{pmatrix} 1 & 0 & \dots & 0 & \lambda_{1,n+1} & \dots & \lambda_{1,m} \\ 0 & 1 & \dots & 0 & \lambda_{2,n+1} & \dots & \lambda_{2,m} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 & \lambda_{n,n+1} & \dots & \lambda_{n,m} \end{pmatrix}$$

satisfying the condition

the column vectors $\lambda_{j_1}, \dots, \lambda_{j_n}$ corresponding to any vertex $v = F_{j_1} \cap \dots \cap F_{j_n}$ form a basis of \mathbb{Z}^n .

We refer to (P, Λ) as a [combinatorial quasitoric pair](#).

Define $K = K(\Lambda) := \ker(\Lambda: T^m \rightarrow T^n) \cong T^{m-n}$.

Prop 2. $K(\Lambda)$ acts freely on \mathcal{Z}_P .

The quotient

$$M = M(P, \Lambda) := \mathcal{Z}_P / K(\Lambda)$$

is the [quasitoric manifold](#) corresponding to (P, Λ) . It has a residual T^n -action ($T^m / K(\Lambda) \cong T^n$) satisfying the two Davis–Januszkiewicz conditions:

- a) the T^n -action is locally standard;
- b) there is a projection $\pi: M \rightarrow P$ whose fibres are orbits of the T^n -action.

Algebraic and symplectic geometers would recognise in the above construction of a quasitoric manifold M from \mathcal{Z}_P a generalisation of the [symplectic reduction](#) construction of a [Hamiltonian toric manifold](#). In the latter case we take $\Lambda = A_P^t$; then M is a toric manifold corresponding to the [Delzant polytope](#)

$$P = \{\mathbf{x} \in \mathbb{R}^n : (\mathbf{a}_i, \mathbf{x}) + b_i \geq 0 \text{ for } 1 \leq i \leq m\}, \quad \mathbf{a}_i \in \mathbb{Z}^n, b_i \in \mathbb{R}.$$

Here we additionally assume the normal vectors \mathbf{a}_i to be *integer*, and the [Delzant condition](#):

for every vertex $v = F_{i_1} \cap \dots \cap F_{i_n}$ of P , the corresponding normal vectors $\mathbf{a}_{i_1}, \dots, \mathbf{a}_{i_n}$ form a basis of \mathbb{Z}^n

to be satisfied.

Then \mathcal{Z}_P is the level set for the [moment map](#) $\mu: \mathbb{C}^m \rightarrow \mathbb{R}^{m-n}$ corresponding to the Hamiltonian action of $K = \text{Ker } \Lambda = \text{Ker } A^t$ on \mathbb{C}^m .

Define complex line bundles

$$\rho_i: \mathbb{Z}_P \times_K \mathbb{C}_i \rightarrow M, \quad 1 \leq i \leq m,$$

where \mathbb{C}_i is the 1-dim complex T^m -representation defined via the quotient projection $\mathbb{C}^m \rightarrow \mathbb{C}_i$ onto the i th factor.

Thm 3 (Davis–Januszkiewicz). *There is an isomorphism of real vector bundles*

$$\tau M \oplus \mathbb{R}^{2(m-n)} \xrightarrow{\cong} \rho_1 \oplus \cdots \oplus \rho_m.$$

This endows M with the **canonical equivariant stably complex structure**. So we may consider its complex cobordism class $[M] \in \Omega_U$.

Thm 4 (Buchstaber-P-Ray). *Every complex cobordism class in $\dim > 2$ contains a quasitoric manifold.*

The complex cobordism ring Ω_U is multiplicatively generated by the cobordism classes $[H_{ij}]$, $0 \leq i \leq j$, of [Milnor hypersurfaces](#)

$$H_{ij} = \{(z_0 : \dots : z_i) \times (w_0 : \dots : w_j) \in \mathbb{C}P^i \times \mathbb{C}P^j : z_0 w_0 + \dots + z_i w_i = 0\}.$$

But H_{ij} is **not** a quasitoric manifold if $i > 1$.

Idea of proof of Thm 4

- 1) Replace each H_{ij} by a quasitoric (in fact, toric) manifold B_{ij} so that $\{B_{ij}\}$ is still a multiplicative generator set for Ω_U . Therefore, every stably complex manifold is cobordant to the disjoint union of products of B_{ij} 's. Every such product is a q-t manifold, but their disjoint union is not.
- 2) Replace disjoint unions by certain connected sums. This is tricky, because you need to take account of both the torus action and the stably complex structure.

- [1] Frédéric Bosio and Laurent Meersseman. *Real quadrics in \mathbb{C}^n , complex manifolds and convex polytopes*. Acta Math. **197** (2006), no. 1, 53–127.
- [2] Victor M Buchstaber and Taras E Panov. *Torus Actions and Their Applications in Topology and Combinatorics*. University Lecture Series, vol. **24**, Amer. Math. Soc., Providence, R.I., 2002.
- [3] Victor M. Buchstaber, Taras E. Panov and Nigel Ray. *Spaces of polytopes and cobordism of quasitoric manifolds*. Moscow Math. J. **7** (2007), no. 2; arXiv:math.AT/0609346.
- [4] Victor M. Buchstaber, Taras E. Panov and Nigel Ray. *Toric genera*. Preprint (2009), to appear in the arXiv soon.
- [5] Michael W. Davis and Tadeusz Januszkiewicz. *Convex polytopes, Coxeter orbifolds and torus actions*. Duke Math. J., **62** (1991), no. 2, 417–451.