

Homotopy theory of moment-angle complexes

based on joint work with Jelena Grbic, Stephen Theriault and Jie Wu

Taras Panov

Lomonosov Moscow State University

Torus Actions: Topology, Geometry and Number Theory
Khabarovsk, 2–7 September 2013

1. Problems

- identifying the homotopy type of the moment-angle complex $\mathcal{Z}_{\mathcal{K}}$ for certain simplicial complexes \mathcal{K} ;
- describing the multiplication and higher Massey products in the Tor-algebra $H^*(\mathcal{Z}_{\mathcal{K}}) = \text{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[\mathcal{K}], \mathbf{k})$ of the face ring $\mathbf{k}[\mathcal{K}]$;
- describing the Yoneda algebra $\text{Ext}_{\mathbf{k}[\mathcal{K}]}(\mathbf{k}, \mathbf{k})$ in terms of generators and relations;
- describing the structure of the Pontryagin algebra $H_*(\Omega DJ(\mathcal{K}))$ and its commutator subalgebra $H_*(\Omega \mathcal{Z}_{\mathcal{K}})$ via iterated and higher Whitehead (Samelson) products;
- identifying the homotopy type of the loop spaces $\Omega DJ(\mathcal{K})$ and $\Omega \mathcal{Z}_{\mathcal{K}}$.

2. Preliminaries

(X, A) a pair of spaces.

\mathcal{K} a simplicial complex on $[m] = \{1, 2, \dots, m\}$, $\emptyset \in \mathcal{K}$.

Given $I = \{i_1, \dots, i_k\} \subset [m]$, set

$$(X, A)^I = Y_1 \times \dots \times Y_m \quad \text{where } Y_i = \begin{cases} X & \text{if } i \in I, \\ A & \text{if } i \notin I. \end{cases}$$

The \mathcal{K} -polyhedral product of (X, A) is

$$(X, A)^{\mathcal{K}} = \bigcup_{I \in \mathcal{K}} (X, A)^I = \bigcup_{I \in \mathcal{K}} \left(\prod_{i \in I} X \times \prod_{i \notin I} A \right).$$

Example

- 1 $(X, A) = (D^2, S^1)$,
 $\mathcal{Z}_{\mathcal{K}} := (D^2, S^1)^{\mathcal{K}}$ the **moment-angle complex**.
It has an action of the torus T^m .
- 2 $(X, A) = (\mathbb{C}P^\infty, pt)$,
 $DJ(\mathcal{K}) := (\mathbb{C}P^\infty, pt)^{\mathcal{K}}$ the **Davis–Januszkiewicz space**.
- 3 $(X, A) = (\mathbb{C}, \mathbb{C}^\times)$,

$$U(\mathcal{K}) := (\mathbb{C}, \mathbb{C}^\times)^{\mathcal{K}} = \mathbb{C}^m \setminus \bigcup_{\{i_1, \dots, i_k\} \notin \mathcal{K}} \{z_{i_1} = \dots = z_{i_k} = 0\}$$

the complement of a **coordinate subspace arrangement**.

Proposition

There exists a deformation retraction

$$\mathcal{Z}_{\mathcal{K}} \hookrightarrow U(\mathcal{K}) \xrightarrow{\simeq} \mathcal{Z}_{\mathcal{K}}$$

There exists a homotopy fibration

$$\begin{array}{ccccc} \mathcal{Z}_{\mathcal{K}} & \longrightarrow & DJ(\mathcal{K}) & \longrightarrow & (\mathbb{C}P^{\infty})^m \\ \parallel & & \parallel & & \parallel \\ (D^2, S^1)^{\mathcal{K}} & & (\mathbb{C}P^{\infty}, pt)^{\mathcal{K}} & & (\mathbb{C}P^{\infty}, \mathbb{C}P^{\infty})^{\mathcal{K}} \end{array}$$

which splits after looping:

$$\Omega DJ(\mathcal{K}) \simeq \Omega \mathcal{Z}_{\mathcal{K}} \times T^m$$

Warning: this is not an H -space splitting

Proposition

There exists an exact sequence of noncommutative algebras

$$1 \longrightarrow H_*(\Omega \mathcal{Z}_{\mathcal{K}}) \longrightarrow H_*(\Omega DJ(\mathcal{K})) \xrightarrow{\text{Ab}} \Lambda[u_1, \dots, u_m] \longrightarrow 1$$

where $\Lambda[u_1, \dots, u_m]$ denotes the exterior algebra and $\deg u_i = 1$.

Let \mathbf{k} denote \mathbb{Z} or a field.

The **face ring** (the **Stanley–Reisner ring**) of \mathcal{K} is given by

$$\mathbf{k}[\mathcal{K}] := \mathbf{k}[v_1, \dots, v_m] / (v_{i_1} \cdots v_{i_k} = 0 \quad \text{if } \{i_1, \dots, i_k\} \notin \mathcal{K})$$

where $\deg v_i = 2$.

Theorem (Buchstaber-P)

$$H^*(DJ(\mathcal{K})) \cong \mathbf{k}[\mathcal{K}]$$

$$H_*(\Omega DJ(\mathcal{K})) \cong \text{Ext}_{\mathbf{k}[\mathcal{K}]}(\mathbf{k}, \mathbf{k}) \quad \mathbf{k} \text{ is a field}$$

$$H^*(\mathcal{Z}_{\mathcal{K}}) \cong \text{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[\mathcal{K}], \mathbf{k})$$

$$\cong H[\Lambda[u_1, \dots, u_m] \otimes \mathbf{k}[\mathcal{K}], d], \quad du_i = v_i, dv_i = 0$$

$$\cong \bigoplus_{I \in \mathcal{K}} \tilde{H}^{*-|I|-1}(\mathcal{K}_I) \quad \mathcal{K}_I = \mathcal{K}|_I$$

$\mathbf{k}[\mathcal{K}]$ is a **Golod ring** if the multiplication and all higher Massey operations in $\text{Tor}_{\mathbf{k}[v_1, \dots, v_m]}(\mathbf{k}[\mathcal{K}], \mathbf{k})$ are trivial.

3. The case of a flag complex

A **missing face** of \mathcal{K} is a subset $I \subset [m]$ such that $I \notin \mathcal{K}$, but every proper subset of I is a simplex of \mathcal{K} .

\mathcal{K} is a **flag complex** if each of its missing faces has two vertices.

Equivalently, \mathcal{K} is flag if any set of vertices of \mathcal{K} which are pairwise connected by edges spans a simplex.

$$\begin{array}{ccc} \{\text{flag complexes on } [m]\} & \xleftrightarrow{1-1} & \{\text{simple graphs on } [m]\} \\ \mathcal{K} & \rightarrow & \mathcal{K}^1 \quad (\text{one-skeleton}) \\ \mathcal{K}(\Gamma) & \leftarrow & \Gamma \end{array}$$

where $\mathcal{K}(\Gamma)$ is the **clique complex** of Γ
(fill in each clique of Γ with a simplex).

A graph Γ is **chordal** if each of its cycles with ≥ 4 edges has a chord.

Equivalently, Γ is chordal if there are no induced cycles of length ≥ 4 .

Theorem (Fulkerson–Gross)

A graph is chordal if and only if its vertices can be ordered in such a way that, for each vertex i , the lesser neighbours of i form a clique.

(**perfect elimination ordering**)

Theorem (Grbic-P-Theriault-Wu)

\mathcal{K} is flag, \mathbf{k} a field. The following are equivalent:


- 1 $\mathbf{k}[\mathcal{K}]$ is a Golod ring;
- 2 the multiplication in $H^*(\mathcal{Z}_{\mathcal{K}})$ is trivial;
- 3 $\Gamma = \mathcal{K}^1$ is a chordal graph;
- 4 $\mathcal{Z}_{\mathcal{K}}$ has homotopy type of a wedge of spheres.

The equivalence (1) \Leftrightarrow (2) \Leftrightarrow (3) was proved by Berglund and Jöllenbeck in 2007.

Implications (1) \Rightarrow (2), (2) \Rightarrow (3) and (4) \Rightarrow (1) are valid for arbitrary \mathcal{K} .

However, (3) \Rightarrow (4) fails in the non-flag case.

Example



The nonzero cohomology groups of $\mathcal{Z}_{\mathcal{K}}$ are

$$H^0 = \mathbb{Z}, \quad H^5 = \mathbb{Z}^{10}, \quad H^6 = \mathbb{Z}^{15}, \quad H^7 = \mathbb{Z}^6 \\ H^9 = \mathbb{Z}/2.$$

All Massey products vanish for dimensional reasons, so \mathcal{K} is Golod (over any field).

Nevertheless, $\mathcal{Z}_{\mathcal{K}}$ is not homotopy equivalent to a wedge of spheres because of the torsion. In fact,

$$\mathcal{Z}_{\mathcal{K}} \simeq (S^5)^{\vee 10} \vee (S^6)^{\vee 15} \vee (S^7)^{\vee 6} \vee \Sigma^7 \mathbb{R}P^2.$$

Question

Assume that $H^*(\mathcal{Z}_{\mathcal{K}})$ has trivial multiplication, so that \mathcal{K} is Golod, over any field. Is it true that $\mathcal{Z}_{\mathcal{K}}$ is a co- H -space, or even a suspension, as in the previous example?

How many spheres in the wedge? What if $\mathcal{Z}_{\mathcal{K}}$ is not a wedge of spheres?

Theorem

For any flag complex \mathcal{K} , there is an isomorphism

$$H_*(\Omega DJ(\mathcal{K})) \cong T\langle u_1, \dots, u_m \rangle / (u_i^2 = 0, u_i u_j + u_j u_i = 0 \text{ for } \{i, j\} \in \mathcal{K})$$

where $T\langle u_1, \dots, u_m \rangle$ is the free algebra on m generators of degree 1.

Remember the exact sequence of non-commutative algebras

$$1 \longrightarrow H_*(\Omega \mathcal{Z}_{\mathcal{K}}) \longrightarrow H_*(\Omega DJ(\mathcal{K})) \xrightarrow{\text{Ab}} \Lambda[u_1, \dots, u_m] \longrightarrow 1$$

Proposition

For any flag complex \mathcal{K} , the Poincaré series of $H_*(\Omega \mathcal{Z}_{\mathcal{K}})$ is given by

$$P(H_*(\Omega \mathcal{Z}_{\mathcal{K}}); t) = \frac{1}{(1+t)^{m-n}(1-h_1 t + \dots + (-1)^n h_n t^n)},$$

where $\mathbf{h}(\mathcal{K}) = (h_0, h_1, \dots, h_n)$ is the h -vector of \mathcal{K} .

Theorem

Assume that \mathcal{K} is flag. The algebra $H_*(\Omega Z_{\mathcal{K}})$, viewed as the commutator subalgebra of $H_*(\Omega DJ(\mathcal{K}))$, is multiplicatively generated by $\sum_{I \subset [m]} \dim \widetilde{H}^0(\mathcal{K}_I)$ iterated commutators of the form

$$[u_j, u_i], \quad [u_{k_1}, [u_j, u_i]], \quad \dots, \quad [u_{k_1}, [u_{k_2}, \dots [u_{k_{m-2}}, [u_j, u_i]] \dots]]$$

where $k_1 < k_2 < \dots < k_p < j > i$, $k_s \neq i$ for any s , and i is the smallest vertex in a connected component not containing j of the subcomplex $\mathcal{K}_{\{k_1, \dots, k_p, j, i\}}$.

Furthermore, this multiplicative generating set is minimal.

Here is an important particular case (corresponding to $\mathcal{K} = m$ points). It is an analogue of the description of a basis in the commutator subalgebra of a free algebra, given by Cohen and Neisendorfer:

Lemma

Let A be the commutator subalgebra of $T\langle u_1, \dots, u_m \rangle / (u_i^2 = 0)$:

$$1 \longrightarrow A \longrightarrow T\langle u_1, \dots, u_m \rangle / (u_i^2 = 0) \longrightarrow \Lambda[u_1, \dots, u_m] \longrightarrow 1$$

where $\deg u_i = 1$.

Then A is a free associative algebra.

It is minimally generated by the iterated commutators of the form

$$[u_j, u_i], \quad [u_{k_1}, [u_j, u_i]], \quad \dots, \quad [u_{k_1}, [u_{k_2}, \dots [u_{k_{m-2}}, [u_j, u_i]] \dots]]$$

where $k_1 < k_2 < \dots < k_p < j > i$ and $k_s \neq i$ for any s .

The number of commutators of length ℓ is equal to $(\ell - 1) \binom{m}{\ell}$.

Corollary

Assume that \mathcal{K} is flag and $\mathcal{Z}_{\mathcal{K}}$ has homotopy type of a wedge of spheres.

Then the number of spheres of dimension $\ell + 1$ in the wedge is given by $\sum_{|I|=\ell} \dim \tilde{H}^0(\mathcal{K}_I)$, for $2 \leq \ell \leq m$.

In particular, $H^i(\mathcal{K}_I) = 0$ for $i > 0$ and all I .

Jelena Grbic, Taras Panov, Stephen Theriault and Jie Wu.
Homotopy types of moment-angle complexes.
Preprint (2012); arXiv:1211.0873.