

Complex geometry of moment-angle manifolds

Taras Panov

joint with Yuri Ustinovsky and Misha Verbitsky

Moscow State University

A Conference to Celebrate Sam Gitler's 80th Birthday
El Colegio Nacional, Mexico City, 25–27 September 2013

1. Moment-angle manifolds from simplicial fans.

Σ a complete simplicial fan in \mathbb{R}^n (not necessarily rational!)

$\mathbf{a}_1, \dots, \mathbf{a}_m \in \mathbb{R}^n$ generators of 1-dimensional cones

$$\mathcal{K} = \mathcal{K}_\Sigma = \left\{ I \subset [m] : \{\mathbf{a}_i : i \in I\} \text{ spans a cone of } \Sigma \right\}$$

the underlying simplicial complex of Σ .

$$\mathcal{Z}_\mathcal{K} = (D^2, S^1)^\mathcal{K} = \bigcup_{I \in \mathcal{K}} \left(\prod_{i \in I} D^2 \times \prod_{i \notin I} S^1 \right) \subset (D^2)^m$$

the **moment-angle manifold** corresponding to \mathcal{K} (or Σ).

$$\begin{aligned} U(\mathcal{K}) &= \mathbb{C}^m \setminus \bigcup_{\{i_1, \dots, i_k\} \notin \mathcal{K}} \{z \in \mathbb{C}^m : z_{i_1} = \dots = z_{i_k} = 0\} \\ &= (\mathbb{C}, \mathbb{C}^\times)^\mathcal{K} = \bigcup_{I \in \mathcal{K}} \left(\prod_{i \in I} \mathbb{C} \times \prod_{i \notin I} \mathbb{C}^\times \right) \end{aligned}$$

the complement of a **coordinate subspace arrangement** corresponding to \mathcal{K} .

Note: $\mathcal{Z}_\mathcal{K}$ is a deformation retract of $U(\mathcal{K})$ for every \mathcal{K} .

Define a map

$$A: \mathbb{R}^m \rightarrow \mathbb{R}^n, \quad \mathbf{e}_i \mapsto \mathbf{a}_i,$$

where $\mathbf{e}_1, \dots, \mathbf{e}_m$ is the standard basis of \mathbb{R}^m . Set

$$\mathbb{R}_{>}^m = \{(y_1, \dots, y_m) \in \mathbb{R}^m : y_i > 0\},$$

and define

$$R := \exp(\text{Ker } A) = \left\{ (y_1, \dots, y_m) \in \mathbb{R}_{>}^m : \prod_{i=1}^m y_i^{\langle \mathbf{a}_i, \mathbf{u} \rangle} = 1 \text{ for all } \mathbf{u} \in \mathbb{R}^n \right\},$$

$R \subset \mathbb{R}_{>}^m$ acts on $U(\mathcal{K}_\Sigma) \subset \mathbb{C}^m$ by coordinatewise multiplications.

Thm 1. *Let Σ be a complete simplicial fan in \mathbb{R}^n with m one-dimensional cones, and let $\mathcal{K} = \mathcal{K}_\Sigma$ be its underlying simplicial complex. Then*

- (a) *the group $R \cong \mathbb{R}^{m-n}$ acts on $U(\mathcal{K})$ freely and properly, so the quotient $U(\mathcal{K})/R$ is a smooth $(m+n)$ -dimensional manifold;*
- (b) *$U(\mathcal{K})/R$ is \mathbb{T}^m -equivariantly homeomorphic to $\mathcal{Z}_\mathcal{K}$.*

Therefore, $\mathcal{Z}_\mathcal{K}$ can be smoothed canonically.

2. Complex-analytic structures.

We shall show that the even-dimensional moment-angle manifold $\mathcal{Z}_{\mathcal{K}}$ corresponding to a complete simplicial fan admits a structure of a complex manifold. The idea is to replace the action of $\mathbb{R}_{>}^{m-n}$ on $U(\mathcal{K})$ (whose quotient is $\mathcal{Z}_{\mathcal{K}}$) by a holomorphic action of $\mathbb{C}^{\frac{m-n}{2}}$ on the same space.

Rem 1. Complex structures on *polytopal* moment-angle manifolds \mathcal{Z}_P were described by [Bosio](#) and [Meersseman](#). They identified \mathcal{Z}_P with a class of complex manifolds known as **LVM-manifolds** (named after [López de Medrano](#), [Verjovsky](#) and [Meersseman](#)).

Topology of polytopal moment-angle manifolds \mathcal{Z}_P is interesting and complicated. [López de Medrano](#) and [Gitler](#) identified their diffeomorphism types for many important series of polytopes.

Assume $m - n$ is even from now on. We can always achieve this by formally adding an ‘empty’ one-dimensional cone to Σ ; this corresponds to adding a ghost vertex to \mathcal{K} , or multiplying $\mathcal{Z}_{\mathcal{K}}$ by a circle.

Set $\ell = \frac{m-n}{2}$.

Constr 1. Choose a linear map $\Psi: \mathbb{C}^\ell \rightarrow \mathbb{C}^m$ satisfying the two conditions:

(a) $\text{Re} \circ \Psi: \mathbb{C}^\ell \rightarrow \mathbb{R}^m$ is a monomorphism.

(b) $A \circ \text{Re} \circ \Psi = 0$.

The composite map of the top line in the following diagram is zero:

$$\begin{array}{ccccccc}
 \mathbb{C}^\ell & \xrightarrow{\Psi} & \mathbb{C}^m & \xrightarrow{\text{Re}} & \mathbb{R}^m & \xrightarrow{A} & \mathbb{R}^n \\
 & & \downarrow \text{exp} & & \downarrow \text{exp} & & \downarrow \text{exp} \\
 & & (\mathbb{C}^\times)^m & \xrightarrow{|\cdot|} & \mathbb{R}_{>}^m & \xrightarrow{\text{exp } A} & \mathbb{R}_{>}^n
 \end{array}$$

where $|\cdot|$ denotes the map $(z_1, \dots, z_m) \mapsto (|z_1|, \dots, |z_m|)$. Now set

$$C = \exp \Psi(\mathbb{C}^\ell) = \left\{ \left(e^{\langle \psi_1, \mathbf{w} \rangle}, \dots, e^{\langle \psi_m, \mathbf{w} \rangle} \right) \in (\mathbb{C}^\times)^m \right\}$$

where $\mathbf{w} = (w_1, \dots, w_\ell) \in \mathbb{C}^\ell$, ψ_i denotes the i th row of the $m \times \ell$ -matrix $\Psi = (\psi_{ij})$.

Then $C \cong \mathbb{C}^\ell$ is a complex-analytic (but not algebraic) subgroup in $(\mathbb{C}^\times)^m$. It acts on $U(\mathcal{K})$ by holomorphic transformations.

Ex 1. Let \mathcal{K} be empty on 2 elements (that is, \mathcal{K} has two ghost vertices). We therefore have $n = 0$, $m = 2$, $\ell = 1$, and $A: \mathbb{R}^2 \rightarrow 0$ is a zero map. Let $\Psi: \mathbb{C} \rightarrow \mathbb{C}^2$ be given by $z \mapsto (z, \alpha z)$ for some $\alpha \in \mathbb{C}$, so that

$$C = \{(e^z, e^{\alpha z})\} \subset (\mathbb{C}^\times)^2.$$

Condition (b) of Constr 1 is void, while (a) is equivalent to that $\alpha \notin \mathbb{R}$. Then $\exp \Psi: \mathbb{C} \rightarrow (\mathbb{C}^\times)^2$ is an embedding, and the quotient $(\mathbb{C}^\times)^2/C$ with the natural complex structure is a complex torus $T_{\mathbb{C}}^2$ with parameter $\alpha \in \mathbb{C}$:

$$(\mathbb{C}^\times)^2/C \cong \mathbb{C}/(\mathbb{Z} \oplus \alpha\mathbb{Z}) = T_{\mathbb{C}}^2(\alpha).$$

Similarly, if \mathcal{K} is empty on 2ℓ elements (so that $n = 0$, $m = 2\ell$), we may obtain any complex torus $T_{\mathbb{C}}^{2\ell}$ as the quotient $(\mathbb{C}^\times)^{2\ell}/C$.

Thm 2. *Let Σ be a complete simplicial fan in \mathbb{R}^n with m one-dimensional cones, and let $\mathcal{K} = \mathcal{K}_\Sigma$ be its underlying simplicial complex. Assume that $m - n = 2\ell$. Then*

- (a) the holomorphic action of the group $C \cong \mathbb{C}^\ell$ on $U(\mathcal{K})$ is free and proper, so the quotient $U(\mathcal{K})/C$ is a compact complex $(m - \ell)$ -manifold;*
- (b) there is a \mathbb{T}^m -equivariant diffeomorphism $U(\mathcal{K})/C \cong \mathcal{Z}_\mathcal{K}$ defining a complex structure on $\mathcal{Z}_\mathcal{K}$ in which \mathbb{T}^m acts holomorphically.*

Ex 2 (Hopf manifold). Let Σ be the complete fan in \mathbb{R}^n whose cones are generated by all proper subsets of $n + 1$ vectors $\mathbf{e}_1, \dots, \mathbf{e}_n, -\mathbf{e}_1 - \dots - \mathbf{e}_n$.

To make $m - n$ even we add one ‘empty’ 1-cone. We have $m = n + 2$, $\ell = 1$. Then $A: \mathbb{R}^{n+2} \rightarrow \mathbb{R}^n$ is given by the matrix $(\mathbf{0} \ I \ -\mathbf{1})$, where I is the unit $n \times n$ matrix, and $\mathbf{0}, \mathbf{1}$ are the n -columns of zeros and units respectively.

We have that \mathcal{K} is the boundary of an n -dim simplex with $n + 1$ vertices and 1 ghost vertex, $\mathcal{Z}_{\mathcal{K}} \cong S^1 \times S^{2n+1}$, and $U(\mathcal{K}) = \mathbb{C}^\times \times (\mathbb{C}^{n+1} \setminus \{0\})$.

Take $\Psi: \mathbb{C} \rightarrow \mathbb{C}^{n+2}$, $z \mapsto (z, \alpha z, \dots, \alpha z)$ for some $\alpha \in \mathbb{C}$, $\alpha \notin \mathbb{R}$. Then

$$C = \{(e^z, e^{\alpha z}, \dots, e^{\alpha z}) : z \in \mathbb{C}\} \subset (\mathbb{C}^\times)^{n+2},$$

and $\mathcal{Z}_{\mathcal{K}}$ acquires a complex structure as the quotient $U(\mathcal{K})/C$:

$$\mathbb{C}^\times \times (\mathbb{C}^{n+1} \setminus \{0\}) / \{(t, \mathbf{w}) \sim (e^z t, e^{\alpha z} \mathbf{w})\} \cong (\mathbb{C}^{n+1} \setminus \{0\}) / \{\mathbf{w} \sim e^{2\pi i \alpha} \mathbf{w}\},$$

where $t \in \mathbb{C}^\times$, $\mathbf{w} \in \mathbb{C}^{n+1} \setminus \{0\}$. The latter quotient of $\mathbb{C}^{n+1} \setminus \{0\}$ is known as the **Hopf manifold**.

3. Holomorphic bundles over toric varieties.

Manifolds $\mathcal{Z}_{\mathcal{K}}$ corresponding to complete *regular* (in particular, *rational*) simplicial fans are total spaces of **holomorphic principal bundles** over **toric varieties** with fibre a complex torus. This allows us to calculate invariants of the complex structures on $\mathcal{Z}_{\mathcal{K}}$, such as Hodge numbers and Dolbeault cohomology.

A **toric variety** is a normal algebraic variety X on which an algebraic torus $(\mathbb{C}^\times)^n$ acts with a dense (Zariski open) orbit.

Toric varieties are classified by *rational* fans. Under this correspondence,

complete fans	\longleftrightarrow	compact varieties
normal fans of polytopes	\longleftrightarrow	projective varieties
regular fans	\longleftrightarrow	nonsingular varieties
simplicial fans	\longleftrightarrow	orbifolds

Σ complete, simplicial, *rational*;

$\mathbf{a}_1, \dots, \mathbf{a}_m$ primitive integral generators of 1-cones;

$\mathbf{a}_i = (a_{i1}, \dots, a_{in}) \in \mathbb{Z}^n$.

Constr 2 ('Cox construction'). Let $A_{\mathbb{C}}: \mathbb{C}^m \rightarrow \mathbb{C}^n$, $\mathbf{e}_i \mapsto \mathbf{a}_i$,

$$\exp A_{\mathbb{C}}: (\mathbb{C}^{\times})^m \rightarrow (\mathbb{C}^{\times})^n,$$

$$(z_1, \dots, z_m) \mapsto \left(\prod_{i=1}^m z_i^{a_{i1}}, \dots, \prod_{i=1}^m z_i^{a_{in}} \right)$$

Set $G = \text{Ker } \exp A_{\mathbb{C}}$.

This is an $(m - n)$ -dimensional algebraic subgroup in $(\mathbb{C}^{\times})^m$.

It acts almost freely (with finite isotropy subgroups) on $U(\mathcal{K}_{\Sigma})$.

If Σ is regular, then $G \cong (\mathbb{C}^{\times})^{m-n}$ and the action is free.

$V_{\Sigma} = U(\mathcal{K}_{\Sigma})/G$ the **toric variety** associated to Σ .

The quotient torus $(\mathbb{C}^{\times})^m/G \cong (\mathbb{C}^{\times})^n$ acts on V_{Σ} with a dense orbit.

Observe that $\mathbb{C}^\ell \cong C \subset G \cong (\mathbb{C}^\times)^{m-n}$ as a complex subgroup.

Prop 1.

- (a) *The toric variety V_Σ is homeomorphic to the quotient of $\mathcal{Z}_{\mathcal{K}_\Sigma}$ by the holomorphic action of G/C .*
- (b) *If Σ is regular, then there is a holomorphic principal bundle $\mathcal{Z}_{\mathcal{K}_\Sigma} \rightarrow V_\Sigma$ with fibre the compact complex torus G/C of dimension ℓ .*

Rem 2. For singular varieties V_Σ the quotient projection $\mathcal{Z}_{\mathcal{K}_\Sigma} \rightarrow V_\Sigma$ is a holomorphic principal **Seifert bundle** for an appropriate orbifold structure on V_Σ .

4. Submanifolds and analytic subsets.

The complex structure on $\mathcal{Z}_{\mathcal{K}}$ is determined by two pieces of data:

- the complete simplicial fan Σ with generators $\mathbf{a}_1, \dots, \mathbf{a}_m$;
- the ℓ -dimensional holomorphic subgroup $C \subset (\mathbb{C}^\times)^m$.

If this data is *generic* (in particular, the fan Σ is not rational), then there is no holomorphic principal torus fibration $\mathcal{Z}_{\mathcal{K}} \rightarrow V_{\Sigma}$ over a toric variety V_{Σ} .

However, there still exists a holomorphic ℓ -dimensional *foliation* \mathcal{F} with a transverse Kähler form $\omega_{\mathcal{F}}$. This form can be used to describe submanifolds and analytic subsets in $\mathcal{Z}_{\mathcal{K}}$.

Consider the complexified map $A_{\mathbb{C}}: \mathbb{C}^m \rightarrow \mathbb{C}^n$, $\mathbf{e}_i \mapsto \mathbf{a}_i$. and the following complex $(m - n)$ -dimensional subgroup in $(\mathbb{C}^\times)^m$:

$$G = \exp(\text{Ker } A_{\mathbb{C}}) = \left\{ \left(e^{z_1}, \dots, e^{z_m} \right) \in (\mathbb{C}^\times)^m : (z_1, \dots, z_m) \in \text{Ker } A_{\mathbb{C}} \right\}.$$

Note $C \subset G$.

The group G acts on $U(\mathcal{K})$, and its orbits define a holomorphic foliation on $U(\mathcal{K})$. Since $G \subset (\mathbb{C}^\times)^m$, this action is free on open subset $(\mathbb{C}^\times)^m \subset U(\mathcal{K})$, so that the generic leaf of the foliation has complex dimension $m - n = 2\ell$.

The ℓ -dimensional closed subgroup $C \subset G$ acts on $U(\mathcal{K})$ freely and properly by Theorem 2, so that $U(\mathcal{K})/C$ carries a holomorphic action of the quotient group $D = G/C$.

\mathcal{F} : the holomorphic foliation on $U(\mathcal{K})/C \cong \mathcal{Z}_{\mathcal{K}}$ by the orbits of D .

The subgroup $G \subset (\mathbb{C}^\times)^m$ is closed if and only if it is isomorphic to $(\mathbb{C}^\times)^{2\ell}$; in this case the subspace $\text{Ker } A \subset \mathbb{R}^m$ is rational. Then Σ is a rational fan and V_Σ is the quotient $U(\mathcal{K})/G$. The foliation \mathcal{F} gives rise to a holomorphic principal Seifert fibration $\pi: \mathcal{Z}_\mathcal{K} \rightarrow V_\Sigma$ with fibres compact complex tori G/C .

For a generic configuration of nonzero vectors $\mathbf{a}_1, \dots, \mathbf{a}_m$, G is biholomorphic to $\mathbb{C}^{2\ell}$ and $D = G/C$ is biholomorphic to \mathbb{C}^ℓ .

A $(1, 1)$ -form $\omega_{\mathcal{F}}$ on the complex manifold $\mathcal{Z}_{\mathcal{K}}$ is called **transverse Kähler** with respect to the foliation \mathcal{F} if

- (a) $\omega_{\mathcal{F}}$ is closed, i.e. $d\omega_{\mathcal{F}} = 0$;
- (b) $\omega_{\mathcal{F}}$ is nonnegative and the zero space of $\omega_{\mathcal{F}}$ is the tangent space of \mathcal{F} .

A complete simplicial fan Σ in \mathbb{R}^n is called **weakly normal** if there exists a (not necessarily simple) n -dimensional polytope P such that Σ is a simplicial subdivision of the normal fan Σ_P .

Thm 3. *Assume that Σ is a weakly normal fan. Then there exists an exact $(1, 1)$ -form $\omega_{\mathcal{F}}$ on $\mathcal{Z}_{\mathcal{K}} = U(\mathcal{K})/C$ which is transverse Kähler for the foliation \mathcal{F} on the dense open subset $(\mathbb{C}^{\times})^m/C \subset U(\mathcal{K})/C$.*

For each $J \subset [m]$, define the corresponding **coordinate submanifold** in $\mathcal{Z}_{\mathcal{K}}$ by

$$\mathcal{Z}_{\mathcal{K}_J} = \{(z_1, \dots, z_m) \in \mathcal{Z}_{\mathcal{K}} : z_i = 0 \text{ for } i \notin J\}.$$

Obviously, $\mathcal{Z}_{\mathcal{K}_J}$ is identified with the quotient of

$$U(\mathcal{K}_J) = \{(z_1, \dots, z_m) \in U(\mathcal{K}) : z_i = 0 \text{ for } i \notin J\}$$

by $C \cong \mathbb{C}^\ell$. In particular, $U(\mathcal{K}_J)/C$ is a complex submanifold in $\mathcal{Z}_{\mathcal{K}} = U(\mathcal{K})/C$.

Observe that the closure of any $(\mathbb{C}^\times)^m$ -orbit of $U(\mathcal{K})$ has the form $U(\mathcal{K}_J)$ for some $J \subset [m]$ (in particular, the dense orbit corresponds to $J = [m]$). Similarly, the closure of any $(\mathbb{C}^\times)^m/C$ -orbit of $\mathcal{Z}_{\mathcal{K}} \cong U(\mathcal{K})/C$ has the form $\mathcal{Z}_{\mathcal{K}_J}$.

Thm 4. *Assume that the data defining a complex structure on $\mathcal{Z}_{\mathcal{K}} = U(\mathcal{K})/C$ is generic. Then any divisor of $\mathcal{Z}_{\mathcal{K}}$ is a union of coordinate divisors.*

Furthermore, if Σ is a weakly normal fan, then any compact irreducible analytic subset $Y \subset \mathcal{Z}_{\mathcal{K}}$ of positive dimension is a coordinate submanifold.

Cor 1. *Under generic assumptions, there are no non-constant meromorphic functions on $\mathcal{Z}_{\mathcal{K}}$.*

- [BP] Victor Buchstaber and Taras Panov. *Torus Actions and Their Applications in Topology and Combinatorics*. University Lecture Series, vol. **24**, Amer. Math. Soc., Providence, R.I., 2002.
- [GL] Samuel Gitler and Santiago López de Medrano. *Intersections of quadrics, moment-angle manifolds and connected sums*. Preprint (2009); arXiv:0901.2580.
- [MR] Laurent Meersseman and Alberto Verjovsky. *Holomorphic principal bundles over projective toric varieties*. J. Reine Angew. Math. **572** (2004), 57–96.
- [PU] Taras Panov and Yuri Ustinovsky. *Complex-analytic structures on moment-angle manifolds*. Moscow Math. J. **12** (2012), no. 1.
- [PUV] Taras Panov, Yuri Ustinovsky and Misha Verbitsky. *Complex geometry of moment-angle manifolds*. Preprint (2013); arXiv:1308.2818.