

Manifolds defined by right-angled 3-dimensional polytopes

Taras Panov

Moscow State University

Princeton-Rider Workshop
on the homotopy theory of polyhedral products

Princeton University, Rider University, 29 May–2 June 2017

Polytopes and moment-angle manifolds

A **convex polytope** in \mathbb{R}^n is a bounded intersection of m halfspaces:

$$P = \{ \mathbf{x} \in \mathbb{R}^n : \langle \mathbf{a}_i, \mathbf{x} \rangle + b_i \geq 0 \text{ for } i = 1, \dots, m \},$$

where $\mathbf{a}_i \in \mathbb{R}^n$ and $b_i \in \mathbb{R}$.

Assume that $F_i = P \cap \{ \mathbf{x} : \langle \mathbf{a}_i, \mathbf{x} \rangle + b_i = 0 \}$ is a facet for each i .
 $\mathcal{F} = \{ F_1, \dots, F_m \}$ the set of facets of P .

Define an affine map

$$i_P: \mathbb{R}^n \rightarrow \mathbb{R}^m, \quad i_P(\mathbf{x}) = (\langle \mathbf{a}_1, \mathbf{x} \rangle + b_1, \dots, \langle \mathbf{a}_m, \mathbf{x} \rangle + b_m).$$

Then i_P is injective, and $i_P(P) \subset \mathbb{R}^m$ is the intersection of an n -dimensional plane with $\mathbb{R}_{\geq}^m = \{ \mathbf{y} = (y_1, \dots, y_m) : y_i \geq 0 \}$.

Define the space \mathcal{Z}_P from the diagram

$$\begin{array}{ccc}
 \mathcal{Z}_P & \xrightarrow{i_Z} & \mathbb{C}^m & & (z_1, \dots, z_m) \\
 \downarrow & & \downarrow \mu & & \downarrow \\
 P & \xrightarrow{i_P} & \mathbb{R}_{\geq}^m & & (|z_1|^2, \dots, |z_m|^2)
 \end{array}$$

Explicitly, $\mathcal{Z}_P = \mu^{-1}(i_P(P))$. It has a T^m -action with the quotient $\mathcal{Z}_P/T^m = P$.

P is **simple** if there are $n = \dim P$ facets meeting at each vertex.

Proposition

If P is a simple polytope, then \mathcal{Z}_P is a smooth $(m+n)$ -dim manifold.

Proof.

Write $i_P(\mathbb{R}^n)$ by $(m-n)$ linear equations in $(y_1, \dots, y_m) \in \mathbb{R}^m$. Replacing each y_k by $|z_k|^2$ we obtain a presentation of \mathcal{Z}_P by Hermitian quadrics. \square

\mathcal{Z}_P is the **moment-angle manifold** (corresponding to P).

Similarly, considering

$$\begin{array}{ccc} \mathcal{R}_P & \longrightarrow & \mathbb{R}^m & (u_1, \dots, u_m) \\ \downarrow & & \downarrow \mu & \downarrow \\ P & \xrightarrow{i_P} & \mathbb{R}_{\geq}^m & (u_1^2, \dots, u_m^2) \end{array}$$

we obtain a **real moment-angle manifold** \mathcal{R}_P .

Example

$P = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 \geq 0, x_2 \geq 0, -\gamma_1 x_1 - \gamma_2 x_2 + 1 \geq 0\}$, $\gamma_1, \gamma_2 > 0$
(a 2-simplex). Then

$\mathcal{Z}_P = \{(z_1, z_2, z_3) \in \mathbb{C}^3 : \gamma_1 |z_1|^2 + \gamma_2 |z_2|^2 + |z_3|^2 = 1\}$ (a 5-sphere),

$\mathcal{R}_P = \{(u_1, u_2, u_3) \in \mathbb{R}^3 : \gamma_1 |u_1|^2 + \gamma_2 |u_2|^2 + |u_3|^2 = 1\}$ (a 2-sphere).

Right-angled polytopes and hyperbolic manifolds

Let P be a polytope in n -dimensional Lobachevsky space \mathbb{L}^n with right angles between adjacent facets (a **right-angled n -polytope**).

Denote by $G(P)$ the group generated by reflections in the facets of P . It is a **right-angled Coxeter group** given by the presentation

$$G(P) = \langle g_1, \dots, g_m \mid g_i^2 = 1, g_i g_j = g_j g_i \text{ if } F_i \cap F_j \neq \emptyset \rangle,$$

where g_i denotes the reflection in the facet F_i .

The group $G(P)$ acts on \mathbb{L}^n discretely with finite isotropy subgroups and with fundamental domain P .

Lemma

Consider an epimorphism $\varphi: G(P) \rightarrow \mathbb{Z}_2^k$, $k \geq n$. The subgroup $\text{Ker } \varphi \subset G(P)$ does not contain elements of finite order if and only if the images of the reflections in any n facets of P that have a common vertex are linearly independent in \mathbb{Z}_2^k .

In this case the group $\text{Ker } \varphi$ acts freely on \mathbb{L}^n .

The quotient $N = \mathbb{L}^n / \text{Ker } \varphi$ is a **hyperbolic n -manifold**. It is composed of $|\mathbb{Z}_2^k| = 2^k$ copies of P and has a Riemannian metric of constant negative curvature. Furthermore, the manifold N is aspherical (the Eilenberg–Mac Lane space $K(\text{Ker } \varphi, 1)$), as its universal cover \mathbb{L}^n is contractible.

Which combinatorial n -polytopes have right-angled realisations in \mathbb{L}^n ?
In dim 3, there is a nice criterion going back to Pogorelov's work of 1967:

Theorem (Pogorelov, Andreev)

A combinatorial 3-polytope $P \neq \Delta^3$ can be realised as a right-angled polytope in \mathbb{L}^3 if and only if it is simple, and does not have 3- and 4-belts of facets. Furthermore, such a realisation is unique up to isometry.

We refer to the above class of 3-polytopes as the **Pogorelov class** \mathcal{P} .
A Pogorelov polytope does not have triangular or quadrangular facets.
The Pogorelov class contains all **fullerenes** (simple 3-polytopes with only pentagonal and hexagonal facets).
The conditions specifying Pogorelov polytopes also feature as the **no- Δ** and **no- \square condition** in Gromov's theory of hyperbolic groups.

There is no classification of right-angled polytopes in \mathbb{L}^4 . For $n \geq 5$, right-angled polytopes in \mathbb{L}^n do not exist [Vinberg].

Given a right-angled polytope P , how to find an epimorphism $\varphi: G(P) \rightarrow \mathbb{Z}_2^k$ with $\text{Ker } \varphi$ acting freely on \mathbb{L}^n ?

One can consider the abelianisation: $G(P) \xrightarrow{\text{ab}} \mathbb{Z}_2^m$, with $\text{Ker ab} = G'(P)$, the **commutator subgroup**.

The corresponding n -manifold $\mathbb{L}^n/G'(P)$ is the real moment-angle manifold \mathcal{R}_P , described as an intersection of quadrics in the beginning of this talk.

Corollary

If P is a right-angled polytope in \mathbb{L}^n , then the real moment-angle manifold \mathcal{R}_P admits a hyperbolic structure as $\mathbb{L}^n/G'(P)$, where $G'(P)$ is the commutator subgroup of the corresponding right-angled Coxeter group. The manifold \mathcal{R}_P is composed of 2^m copies of P .

A more economical way to obtain a hyperbolic manifold is to consider $\varphi: G(P) \rightarrow \mathbb{Z}_2^n$. Such an epimorphism factors as $G(P) \xrightarrow{\text{ab}} \mathbb{Z}_2^m \xrightarrow{\Lambda} \mathbb{Z}_2^n$, where Λ is a linear map.

The subgroup $\text{Ker } \varphi$ acts freely on \mathbb{L}^n if and only if the Λ -images of any n facets of P that meet at a vertex form a basis of \mathbb{Z}_2^n . Such Λ is called a **\mathbb{Z}_2 -characteristic function**.

Proposition

Any simple 3-polytope admits a characteristic function.

Proof.

Given a 4-colouring of the facets of P , we assign to a facet of i th colour the i th basis vector $\mathbf{e}_i \in \mathbb{Z}^3$ for $i = 1, 2, 3$ and the vector $\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3$ for $i = 4$. The resulting map $\Lambda: \mathbb{Z}_2^m \rightarrow \mathbb{Z}_2^3$ satisfies the required condition, as any three of the four vectors $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3$ form a basis of \mathbb{Z}^3 . \square

Definition (A. Vesnin, 1987)

$N(P, \Lambda) = \mathbb{L}^3 / \text{Ker } \varphi$ a **hyperbolic 3-manifold of Löbell type**.

It is composed of $|\mathbb{Z}_2^3| = 8$ copies of a right-angled 3-polytope $P \in \mathcal{P}$ glued along their facets;

the gluing is prescribed by the characteristic function $\Lambda: \mathbb{Z}_2^m \rightarrow \mathbb{Z}_2^3$.



In particular, one obtains a hyperbolic 3-manifold $N(P, \chi)$ from any regular 4-colouring $\chi: \mathcal{F} \rightarrow \{1, 2, 3, 4\}$ of a right-angled 3-polytope P .

Löbell (1931) was first to consider hyperbolic 3-manifolds coming from 4-colourings of a family of right-angled polytopes starting from the dodecahedron.

Example: Löbell polytopes Q_k (“barrel” fullerenes)


For $k \geq 5$, let Q_k be a simple 3-polytope with two “top” and “bottom” k -gonal facets and $2k$ pentagonal facets forming two k -belts around the top and bottom, so that Q_k has $2k + 2$ facets in total.

Note that Q_5 is a combinatorial dodecahedron, while Q_6 is a fullerene.



It is easy to see that $Q_k \in \mathcal{P}$, so it admits a right-angled realisation in \mathbb{L}^3 .

Consider hyperbolic 3-manifolds $N(Q_k, \chi)$ corresponding to 4-colourings χ of Q_k . For example, a dodecahedron Q_5 has a unique 4-colouring up to equivalence, while Q_6 has four non-equivalent regular 4-colourings (4-colourings are equivalent if they differ by a permutation of colours).



Conjecture (Vesnin, 1991)

Hyperbolic 3-manifolds $N(Q_k, \chi_1)$ and $N(Q_k, \chi_2)$ are diffeomorphic (isometric) if and only if the 4-colourings χ_1 and χ_2 are equivalent.

By 2009 the conjecture was verified for all k except 6, 8 using deep results on arithmetic groups (**Margulis commensurator theorem**).

However, it remained open for Q_6 and Q_8 .

Pairs (P, Λ) and (P', Λ') are **equivalent** if P and P' are combinatorially equivalent, and $\Lambda, \Lambda': \mathbb{Z}_2^m \rightarrow \mathbb{Z}_2^n$ differ by an automorphism of \mathbb{Z}_2^n .

Theorem (Buchstaber–Erokhovets–Masuda–P–Park)

Let $N = N(P, \Lambda)$ and $N' = N(P', \Lambda')$ be two hyperbolic 3-manifolds of Löbell type corresponding to right-angled 3-polytopes P and P' . Then the following conditions are equivalent:

(a) there is a cohomology ring isomorphism

$$\varphi: H^*(N; \mathbb{Z}_2) \xrightarrow{\cong} H^*(N'; \mathbb{Z}_2);$$

(b) there is a diffeomorphism $N \cong N'$;

(c) there is an equivalence of \mathbb{Z}_2 -characteristic pairs $(P, \Lambda) \sim (P', \Lambda')$.

The difficult implication is $(a) \Rightarrow (c)$. Its proof builds upon the wealth of cohomological techniques of toric topology.

Specifying to \mathbb{Z}_2 -characteristic functions Λ coming from colourings χ we obtain:

Theorem (Buchstaber–P)

Hyperbolic 3-manifolds $N(P, \chi_1)$ and $N(P', \chi_2)$ corresponding to right-angled polytopes P and P' are diffeomorphic (isometric) if and only if the 4-colourings χ_1 and χ_2 are equivalent.

In particular, Vesnin's conjecture holds for *all* Löbell polytopes Q_k .

Cohomology of moment-angle manifolds

The **face ring** (the **Stanley–Reisner ring**) of a simple polytope P

$$\mathbb{Z}[P] := \mathbb{Z}[v_1, \dots, v_m] / (v_{i_1} \cdots v_{i_k} = 0 \text{ for } F_{i_1} \cap \cdots \cap F_{i_k} = \emptyset)$$

where $\deg v_i = 2$.

Theorem

There are ring isomorphisms

$$\begin{aligned} H^*(\mathcal{Z}_P) &\cong \mathrm{Tor}_{\mathbb{Z}[v_1, \dots, v_m]}(\mathbb{Z}[P], \mathbb{Z}) \\ &\cong H(\Lambda[u_1, \dots, u_m] \otimes \mathbb{Z}[P], d) && du_i = v_i, dv_i = 0 \\ &\cong \bigoplus_{I \subset \{1, \dots, m\}} \tilde{H}^{*-|I|-1}(P_I) && P_I = \bigcup_{i \in I} F_i \end{aligned}$$

(Quasi)toric manifolds and small covers

P a simple n -polytope, $\mathcal{F} = \{F_1, \dots, F_m\}$ the set of facets.

A **characteristic function** is map $\Lambda: \mathcal{F} \rightarrow \mathbb{Z}^n$ such that $\Lambda(F_{i_1}), \dots, \Lambda(F_{i_n})$ is a basis of \mathbb{Z}^n whenever F_{i_1}, \dots, F_{i_n} intersect in a vertex.

A characteristic function defines a linear map $\Lambda: \mathbb{Z}^{\mathcal{F}} = \mathbb{Z}^m \rightarrow \mathbb{Z}^n$ and a homomorphism of tori $\Lambda: T^m \rightarrow T^n$.

Proposition

The subgroup $\text{Ker } \Lambda \cong T^{m-n}$ acts freely on \mathcal{Z}_P .

$M(P, \Lambda) = \mathcal{Z}_P / \text{Ker } \Lambda$ is a **quasitoric manifold**.

It is a smooth $2n$ -dimensional manifold with an action of the n -torus $T^m / \text{Ker } \Lambda \cong T^n$ and the quotient P .

By considering \mathbb{Z}_2 -characteristic functions $\Lambda: \mathbb{Z}_2^m \rightarrow \mathbb{Z}_2^n$, we obtain **small covers** of P as the quotients $\mathcal{R}_P / \text{Ker } \Lambda$.

A small cover $N(P, \Lambda)$ is a smooth n -dimensional manifold with an action of \mathbb{Z}_2^n and the quotient P .

Proposition

In dimension 3, a real moment-angle manifold \mathcal{R}_P and a small cover $N(P, \Lambda)$ admit a hyperbolic structure if and only if P is a Pogorelov polytope.

Proof.

P is a Pogorelov polytope \Leftrightarrow dual \mathcal{K} has no \triangle and \square

If P is a Pogorelov polytope, then \mathcal{R}_P and $N(P, \Lambda)$ are hyperbolic.

If \mathcal{K} has a \triangle , then $R_\triangle \cong S^2$ retracts off \mathcal{R}_P , so \mathcal{R}_P cannot be hyperbolic (as it has $\pi_2 \neq 0$).

If \mathcal{K} has a \square , then $R_\square \cong T^2$ retracts off \mathcal{R}_P , which is impossible for a hyperbolic manifold. □

Theorem (Danilov–Jurkiewicz, Davis–Januszkiewicz)

Let $M = M(P, \Lambda)$ be a quasitoric manifold over a simple n -polytope P . The cohomology ring $H^*(M; \mathbb{Z})$ is generated by the 2-dimensional classes $[v_i]$ dual to the characteristic submanifolds M_i , $i = 1, \dots, m$, and is given by

$$H^*(M; \mathbb{Z}) \cong \mathbb{Z}[v_1, \dots, v_m]/\mathcal{I}, \quad \deg v_i = 2,$$

where \mathcal{I} is the ideal generated by elements of two kinds:

- (a) $v_{i_1} \cdots v_{i_k}$, where $F_{i_1} \cap \cdots \cap F_{i_k} = \emptyset$ in P ;
- (b) $\sum_{i=1}^m \langle \Lambda(F_i), \mathbf{x} \rangle v_i$ for any $\mathbf{x} \in \mathbb{Z}^n$.

The \mathbb{Z}_2 -cohomology ring $H^*(N; \mathbb{Z}_2)$ of a small cover $N = N(P, \Lambda)$ has the same description, with generators v_i of degree 1.

Theorem (Buchstaber-Erokhovets-Masuda-P-Park)

Let $M = M(P, \Lambda)$ and $M' = M(P', \Lambda')$ be quasitoric 6-manifolds, where P is a Pogorelov 3-polytope. The following conditions are equivalent:

- (a) There is a ring isomorphism $\varphi: H^*(M; \mathbb{Z}) \xrightarrow{\cong} H^*(M'; \mathbb{Z})$;
- (b) There is a diffeomorphism $M \cong M'$;
- (c) There is an equivalence of characteristic pairs $(P, \Lambda) \sim (P', \Lambda')$.

Idea of proof (for both theorems).

We need to prove that a ring iso $\varphi: H^*(N; \mathbb{Z}) \xrightarrow{\cong} H^*(N'; \mathbb{Z})$ implies an equivalence $(P, \Lambda) \sim (P', \Lambda')$.

An iso $\varphi: H^*(N; \mathbb{Z}_2) \xrightarrow{\cong} H^*(N'; \mathbb{Z}_2)$ implies an iso

$\varphi: H^*(M; \mathbb{Z}_2) \xrightarrow{\cong} H^*(M'; \mathbb{Z}_2)$ (as every \mathbb{Z}_2 -characteristic function of a 3-polytope lifts to a \mathbb{Z} -characteristic function!)

An iso $\varphi: H^*(M; \mathbb{Z}_2) = \mathbb{Z}_2[P]/\mathcal{J}_\Lambda \xrightarrow{\cong} \mathbb{Z}_2[P']/\mathcal{J}_{\Lambda'} = H^*(M'; \mathbb{Z}_2)$ implies an iso

$$\begin{aligned} \psi: H^*(\mathcal{Z}_P; \mathbb{Z}_2) &= \mathrm{Tor}_{\mathbb{Z}_2[v_1, \dots, v_m]/\mathcal{J}_\Lambda}(\mathbb{Z}_2[P]/\mathcal{J}_\Lambda, \mathbb{Z}_2) \\ &\xrightarrow{\cong} \mathrm{Tor}_{\mathbb{Z}_2[v_1, \dots, v_m]/\mathcal{J}_{\Lambda'}}(\mathbb{Z}_2[P']/\mathcal{J}_{\Lambda'}, \mathbb{Z}_2) = H^*(\mathcal{Z}_{P'}; \mathbb{Z}_2) \end{aligned}$$

The results of [Fan](#), [Ma](#) and [Wang](#) imply that ψ maps the set of canonical generators $\{[u_i v_j] \in H^3(\mathcal{Z}_P): F_i \cap F_j = \emptyset\}$ bijectively to the corresponding set for $\mathcal{Z}_{P'}$.

This implies that φ maps the set $\{[v_i] \in H^2(M)\}$ bijectively to the corresponding set for M' , giving an equivalence $(P, \Lambda) \sim (P', \Lambda')$. □

Problem

Let $M = M(P, \Lambda)$ and $M' = M(P', \Lambda')$ be quasitoric manifolds with isomorphic integer cohomology rings. Are they homeomorphic?

Our result gives a positive answer in the case of quasitoric 6-manifolds over Pogorelov polytopes.

Proposition

6-dimensional quasitoric manifolds M and M' are diffeomorphic if there is an isomorphism $\varphi: H^*(M; \mathbb{Z}) \xrightarrow{\cong} H^*(M'; \mathbb{Z})$ preserving the first Pontryagin class, i. e. $\varphi(p_1(M)) = p_1(M')$.

We have $p_1(M) = v_1^2 + \dots + v_m^2 \in H^4(M)$. However, we were not able to establish the invariance of $p_1(M)$ directly...

- [1] V. Buchstaber, N. Erokhovets, M. Masuda, T. Panov and S. Park. *Cohomological rigidity of manifolds defined by right-angled 3-dimensional polytopes*. Russian Math. Surveys 72 (2017), no. 2; arXiv:1610.07575.
- [2] V. Buchstaber and T. Panov. *On manifolds defined by 4-colourings of simple 3-polytopes*. Russian Math. Surveys 71 (2016), no. 6, 1137-1139; arXiv:1703.06801.