

# Right-angled polytopes, hyperbolic manifolds and torus actions

Taras Panov

Moscow State University

International Conference  
Morse Theory and its Applications  
dedicated to the 70th anniversary of V. Sharko

Kiev, Ukraine  
25–28 September 2019

# Polytopes and moment-angle manifolds

A **convex polytope** in  $\mathbb{R}^n$  is a bounded intersection of  $m$  halfspaces:

$$P = \{ \mathbf{x} \in \mathbb{R}^n : \langle \mathbf{a}_i, \mathbf{x} \rangle + b_i \geq 0 \text{ for } i = 1, \dots, m \},$$

where  $\mathbf{a}_i \in \mathbb{R}^n$  and  $b_i \in \mathbb{R}$ .

Assume that  $F_i = P \cap \{ \mathbf{x} : \langle \mathbf{a}_i, \mathbf{x} \rangle + b_i = 0 \}$  is a facet for each  $i$ .  
 $\mathcal{F} = \{ F_1, \dots, F_m \}$  the set of facets of  $P$ .

Define an affine map

$$i_P: \mathbb{R}^n \rightarrow \mathbb{R}^m, \quad i_P(\mathbf{x}) = (\langle \mathbf{a}_1, \mathbf{x} \rangle + b_1, \dots, \langle \mathbf{a}_m, \mathbf{x} \rangle + b_m).$$

Then  $i_P$  is injective, and  $i_P(P) \subset \mathbb{R}^m$  is the intersection of an  $n$ -dimensional plane with  $\mathbb{R}_{\geq}^m = \{ \mathbf{y} = (y_1, \dots, y_m) : y_i \geq 0 \}$ .

Define the space  $\mathcal{Z}_P$  from the diagram

$$\begin{array}{ccc}
 \mathcal{Z}_P & \xrightarrow{i_Z} & \mathbb{C}^m & (z_1, \dots, z_m) \\
 \downarrow & & \downarrow \mu & \downarrow \\
 P & \xrightarrow{i_P} & \mathbb{R}_{\geq}^m & (|z_1|^2, \dots, |z_m|^2)
 \end{array}$$

Explicitly,  $\mathcal{Z}_P = \mu^{-1}(i_P(P))$ . It has a  $T^m$ -action with the quotient  $\mathcal{Z}_P/T^m = P$ .

$P$  is **simple** if there are  $n = \dim P$  facets meeting at each vertex.

## Proposition

*If  $P$  is a simple polytope, then  $\mathcal{Z}_P$  is a smooth  $(m+n)$ -dim manifold.*

## Proof.

Write  $i_P(\mathbb{R}^n)$  by  $(m-n)$  linear equations in  $(y_1, \dots, y_m) \in \mathbb{R}^m$ . Replacing each  $y_k$  by  $|z_k|^2$  we obtain a presentation of  $\mathcal{Z}_P$  by Hermitian quadrics.  $\square$

$\mathcal{Z}_P$  is the **moment-angle manifold** (corresponding to  $P$ ).

Similarly, considering

$$\begin{array}{ccc} \mathcal{R}_P & \longrightarrow & \mathbb{R}^m & (u_1, \dots, u_m) \\ \downarrow & & \downarrow \mu & \downarrow \\ P & \xrightarrow{i_P} & \mathbb{R}_{\geq}^m & (u_1^2, \dots, u_m^2) \end{array}$$

we obtain a **real moment-angle manifold**  $\mathcal{R}_P$ .

### Example

$P = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 \geq 0, x_2 \geq 0, -\gamma_1 x_1 - \gamma_2 x_2 + 1 \geq 0\}$ ,  $\gamma_1, \gamma_2 > 0$   
(a 2-simplex). Then

$\mathcal{Z}_P = \{(z_1, z_2, z_3) \in \mathbb{C}^3 : \gamma_1 |z_1|^2 + \gamma_2 |z_2|^2 + |z_3|^2 = 1\}$  (a 5-sphere),

$\mathcal{R}_P = \{(u_1, u_2, u_3) \in \mathbb{R}^3 : \gamma_1 |u_1|^2 + \gamma_2 |u_2|^2 + |u_3|^2 = 1\}$  (a 2-sphere).

# Right-angled polytopes and hyperbolic manifolds

Let  $P$  be a polytope in  $n$ -dimensional Lobachevsky space  $\mathbb{L}^n$  with right angles between adjacent facets (a **right-angled  $n$ -polytope**).

Denote by  $G(P)$  the group generated by reflections in the facets of  $P$ . It is a **right-angled Coxeter group** given by the presentation

$$G(P) = \langle g_1, \dots, g_m \mid g_i^2 = 1, g_i g_j = g_j g_i \text{ if } F_i \cap F_j \neq \emptyset \rangle,$$

where  $g_i$  denotes the reflection in the facet  $F_i$ .

The group  $G(P)$  acts on  $\mathbb{L}^n$  discretely with finite isotropy subgroups and with fundamental domain  $P$ .

### Lemma (A. Vesnin, 1987)

*Consider an epimorphism  $\varphi: G(P) \rightarrow \mathbb{Z}_2^k$ ,  $k \geq n$ . The subgroup  $\text{Ker } \varphi \subset G(P)$  does not contain elements of finite order if and only if the images of the reflections in any  $n$  facets of  $P$  that have a common vertex are linearly independent in  $\mathbb{Z}_2^k$ .*

*In this case the group  $\text{Ker } \varphi$  acts freely on  $\mathbb{L}^n$ .*

The quotient  $N = \mathbb{L}^n / \text{Ker } \varphi$  is a **hyperbolic  $n$ -manifold**. It is composed of  $|\mathbb{Z}_2^k| = 2^k$  copies of  $P$  and has a Riemannian metric of constant negative curvature. Furthermore, the manifold  $N$  is aspherical (the Eilenberg–Mac Lane space  $K(\text{Ker } \varphi, 1)$ ), as its universal cover  $\mathbb{L}^n$  is contractible.

Which combinatorial  $n$ -polytopes have right-angled realisations in  $\mathbb{L}^n$ ?  
In dim 3, there is a nice criterion going back to Pogorelov's work of 1967:



Alexei Pogorelov (1919–2002)

## МАТЕМАТИЧЕСКИЕ ЗАМЕТКИ

т. 1, № 1 [1967], 3—8

### О ПРАВИЛЬНОМ РАЗБИЕНИИ ПРОСТРАНСТВА ЛОБАЧЕВСКОГО

А. В. Погорелов

Пусть  $P$  — замкнутый выпуклый многогранник с прямыми двугранными углами в пространстве Лобачевского. Отражим многогранник  $P$  зеркально в каждой из его граней. Полученные при этом многогранники отразим в гранях, не примыкающих к  $P$ , и т. д. Продолжая этот процесс неограниченно, мы заполним все пространство многогранниками, равными  $P$ . Построенное разбиение пространства Лобачевского на равные многогранники осуществляется системой плоскостей. Каждые две плоскости этой системы либо не пересекаются, либо пересекаются под прямым углом.

Рассмотрим пример. На плоскости Лобачевского можно построить правильный пятиугольник с любыми внутренними углами, меньшими  $\frac{3\pi}{5}$ . В частности, существует правильный пятиугольник с прямыми внутренними углами. Из таких плоских пятиугольников, очевидно, составляется правильный додекаэдр в пространстве Лобачевского. Все его двугранные углы прямые, и, следовательно, такими додекаэдрами покрывается все пространство Лобачевского указанным способом. Можно было бы привести и другие примеры многогранников, которыми заполняется пространство Лобачевского.

Целью настоящей заметки является установление необходимых и достаточных условий, при которых существует замкнутый выпуклый многогранник заданного строения с прямыми двугранными углами. При этом

## Theorem (Pogorelov, Andreev)

*A combinatorial 3-polytope  $P \neq \Delta^3$  can be realised as a right-angled polytope in  $\mathbb{L}^3$  if and only if it is simple, and does not have 3- and 4-belts of facets. Furthermore, such a realisation is unique up to isometry.*

We refer to the class of combinatorial 3-polytope satisfying the conditions above as a **Pogorelov class**  $\mathcal{P}$ .

A Pogorelov polytope does not have triangular or quadrangular facets.

Every **fullerene** (a simple 3-polytope with only pentagonal and hexagonal facets) is a Pogorelov polytope.

The conditions specifying Pogorelov polytopes also feature as the **no- $\Delta$**  and **no- $\square$  condition** in Gromov's theory of hyperbolic groups.

Classification of right-angled polytopes in  $\mathbb{L}^4$  is wide open.

There are no right-angled polytopes in  $\mathbb{L}^n$  for  $n \geq 5$ , [Nikulin, Vinberg].



Given a right-angled polytope  $P$ , how to find an epimorphism  $\varphi: G(P) \rightarrow \mathbb{Z}_2^k$  with  $\text{Ker } \varphi$  acting freely on  $\mathbb{L}^n$ ?

One can consider the abelianisation:  $G(P) \xrightarrow{\text{ab}} \mathbb{Z}_2^m$ , with  $\text{Ker ab} = G'(P)$ , the **commutator subgroup**.

The corresponding  $n$ -manifold  $\mathbb{L}^n/G'(P)$  is the real moment-angle manifold  $\mathcal{R}_P$ , described as an intersection of quadrics in the beginning of this talk.

## Corollary

*If  $P$  is a right-angled polytope in  $\mathbb{L}^n$ , then the real moment-angle manifold  $\mathcal{R}_P$  admits a hyperbolic structure as  $\mathbb{L}^n/G'(P)$ , where  $G'(P)$  is the commutator subgroup of the corresponding right-angled Coxeter group. The manifold  $\mathcal{R}_P$  is composed of  $2^m$  copies of  $P$ .*

A more economical way to obtain a hyperbolic manifold is to consider  $\varphi: G(P) \rightarrow \mathbb{Z}_2^n$ . Such an epimorphism factors as  $G(P) \xrightarrow{\text{ab}} \mathbb{Z}_2^m \xrightarrow{\Lambda} \mathbb{Z}_2^n$ , where  $\Lambda$  is a linear map.

The subgroup  $\text{Ker } \varphi$  acts freely on  $\mathbb{L}^n$  if and only if the  $\Lambda$ -images of any  $n$  facets of  $P$  that meet at a vertex form a basis of  $\mathbb{Z}_2^n$ . Such  $\Lambda$  is called a  **$\mathbb{Z}_2$ -characteristic function**.

## Proposition

*Any simple 3-polytope admits a characteristic function.*

## Proof.

Given a 4-colouring of the facets of  $P$ , we assign to a facet of  $i$ th colour the  $i$ th basis vector  $\mathbf{e}_i \in \mathbb{Z}^3$  for  $i = 1, 2, 3$  and the vector  $\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3$  for  $i = 4$ . The resulting map  $\Lambda: \mathbb{Z}_2^m \rightarrow \mathbb{Z}_2^3$  satisfies the required condition, as any three of the four vectors  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3$  form a basis of  $\mathbb{Z}^3$ .  $\square$

## Definition (A. Vesnin, 1987)

$N(P, \Lambda) = \mathbb{L}^3 / \text{Ker } \varphi$  a **hyperbolic 3-manifold of Löbell type**.

It is composed of  $|\mathbb{Z}_2^3| = 8$  copies of a right-angled 3-polytope  $P \in \mathcal{P}$  glued along their facets;

the gluing is prescribed by the characteristic function  $\Lambda: \mathbb{Z}_2^m \rightarrow \mathbb{Z}_2^3$ .



In particular, one obtains a hyperbolic 3-manifold  $N(P, \chi)$  from any regular 4-colouring  $\chi: \mathcal{F} \rightarrow \{1, 2, 3, 4\}$  of a right-angled 3-polytope  $P$ .

Löbell (1931) was first to consider hyperbolic 3-manifolds coming from 4-colourings of a family of right-angled polytopes starting from the dodecahedron.

## Example: Löbell polytopes $Q_k$ (“barrel” fullerenes)


For  $k \geq 5$ , let  $Q_k$  be a simple 3-polytope with two “top” and “bottom”  $k$ -gonal facets and  $2k$  pentagonal facets forming two  $k$ -belts around the top and bottom, so that  $Q_k$  has  $2k + 2$  facets in total.

Note that  $Q_5$  is a combinatorial dodecahedron, while  $Q_6$  is a fullerene.



It is easy to see that  $Q_k \in \mathcal{P}$ , so it admits a right-angled realisation in  $\mathbb{L}^3$ .

Consider hyperbolic 3-manifolds  $N(Q_k, \chi)$  corresponding to 4-colourings  $\chi$  of  $Q_k$ . For example, a dodecahedron  $Q_5$  has a unique 4-colouring up to equivalence, while  $Q_6$  has four non-equivalent regular 4-colourings (4-colourings are equivalent if they differ by a permutation of colours).



### Conjecture (Vesnin, 1991)

*Hyperbolic 3-manifolds  $N(Q_k, \chi_1)$  and  $N(Q_k, \chi_2)$  are diffeomorphic (isometric) if and only if the 4-colourings  $\chi_1$  and  $\chi_2$  are equivalent.*

By 2009 the conjecture was verified for all  $k$  except 6, 8 using deep results on arithmetic groups (**Margulis commensurator theorem**).

However, it remained open for  $Q_6$  and  $Q_8$ .

Pairs  $(P, \Lambda)$  and  $(P', \Lambda')$  are **equivalent** if  $P$  and  $P'$  are combinatorially equivalent, and  $\Lambda, \Lambda': \mathbb{Z}_2^m \rightarrow \mathbb{Z}_2^n$  differ by an automorphism of  $\mathbb{Z}_2^n$ .

### Theorem (Buchstaber–Erokhovets–Masuda–P–Park)

Let  $N = N(P, \Lambda)$  and  $N' = N(P', \Lambda')$  be two hyperbolic 3-manifolds of Löbell type corresponding to right-angled 3-polytopes  $P$  and  $P'$ . Then the following conditions are equivalent:

(a) there is a cohomology ring isomorphism

$$\varphi: H^*(N; \mathbb{Z}_2) \xrightarrow{\cong} H^*(N'; \mathbb{Z}_2);$$

(b) there is a diffeomorphism  $N \cong N'$ ;

(c) there is an equivalence of  $\mathbb{Z}_2$ -characteristic pairs  $(P, \Lambda) \sim (P', \Lambda')$ .

The difficult implication is  $(a) \Rightarrow (c)$ . Its proof builds upon the wealth of cohomological techniques of toric topology.

Specifying to  $\mathbb{Z}_2$ -characteristic functions  $\Lambda$  coming from colourings  $\chi$  we obtain:

### Theorem (Buchstaber–P)

*Hyperbolic 3-manifolds  $N(P, \chi_1)$  and  $N(P', \chi_2)$  corresponding to right-angled polytopes  $P$  and  $P'$  are diffeomorphic (isometric) if and only if the 4-colourings  $\chi_1$  and  $\chi_2$  are equivalent.*

In particular, Vesnin's conjecture holds for all Löbell polytopes  $Q_k$ .

# Cohomology of moment-angle manifolds

The **face ring** (the **Stanley–Reisner ring**) of a simple polytope  $P$

$$\mathbb{Z}[P] := \mathbb{Z}[v_1, \dots, v_m] / (v_{i_1} \cdots v_{i_k} = 0 \text{ for } F_{i_1} \cap \cdots \cap F_{i_k} = \emptyset)$$

where  $\deg v_i = 2$ .

## Theorem

*There are ring isomorphisms*

$$\begin{aligned} H^*(\mathcal{Z}_P) &\cong \text{Tor}_{\mathbb{Z}[v_1, \dots, v_m]}(\mathbb{Z}[P], \mathbb{Z}) \\ &\cong H(\Lambda[u_1, \dots, u_m] \otimes \mathbb{Z}[P], d) && du_i = v_i, dv_i = 0 \\ &\cong \bigoplus_{I \subset \{1, \dots, m\}} \tilde{H}^{*-|I|-1}(P_I) && P_I = \bigcup_{i \in I} F_i \end{aligned}$$



# (Quasi)toric manifolds and small covers

$P$  a simple  $n$ -polytope,  $\mathcal{F} = \{F_1, \dots, F_m\}$  the set of facets.

A **characteristic function** is map  $\Lambda: \mathcal{F} \rightarrow \mathbb{Z}^n$  such that  $\Lambda(F_{i_1}), \dots, \Lambda(F_{i_n})$  is a basis of  $\mathbb{Z}^n$  whenever  $F_{i_1}, \dots, F_{i_n}$  intersect in a vertex.

A characteristic function defines a linear map  $\Lambda: \mathbb{Z}^{\mathcal{F}} = \mathbb{Z}^m \rightarrow \mathbb{Z}^n$  and a homomorphism of tori  $\Lambda: T^m \rightarrow T^n$ .

## Proposition

*The subgroup  $\text{Ker } \Lambda \cong T^{m-n}$  acts freely on  $\mathcal{Z}_P$ .*

$M(P, \Lambda) = \mathcal{Z}_P / \text{Ker } \Lambda$  is a **quasitoric manifold**.

It is a smooth  $2n$ -dimensional manifold with an action of the  $n$ -torus  $T^m / \text{Ker } \Lambda \cong T^n$  and the quotient  $P$ .

By considering  $\mathbb{Z}_2$ -characteristic functions  $\Lambda: \mathbb{Z}_2^m \rightarrow \mathbb{Z}_2^n$ , we obtain **small covers** of  $P$  as the quotients  $\mathcal{R}_P / \text{Ker } \Lambda$ .

A small cover  $N(P, \Lambda)$  is a smooth  $n$ -dimensional manifold with an action of  $\mathbb{Z}_2^n$  and the quotient  $P$ .

## Proposition

*In dimension 3, a real moment-angle manifold  $\mathcal{R}_P$  and a small cover  $N(P, \Lambda)$  admit a hyperbolic structure if and only if  $P$  is a Pogorelov polytope.*

## Proof.

$P$  is a Pogorelov polytope  $\Leftrightarrow$  dual  $\mathcal{K}$  has no  $\triangle$  and  $\square$

If  $P$  is a Pogorelov polytope, then  $\mathcal{R}_P$  and  $N(P, \Lambda)$  are hyperbolic.

If  $\mathcal{K}$  has a  $\triangle$ , then  $R_\triangle \cong S^2$  retracts off  $\mathcal{R}_P$ , so  $\mathcal{R}_P$  cannot be hyperbolic (as it has  $\pi_2 \neq 0$ ).

If  $\mathcal{K}$  has a  $\square$ , then  $R_\square \cong T^2$  retracts off  $\mathcal{R}_P$ , which is impossible for a hyperbolic manifold. □

## Theorem (Danilov–Jurkiewicz, Davis–Januszkiewicz)

Let  $M = M(P, \Lambda)$  be a quasitoric manifold over a simple  $n$ -polytope  $P$ . The cohomology ring  $H^*(M; \mathbb{Z})$  is generated by the 2-dimensional classes  $[v_i]$  dual to the characteristic submanifolds  $M_i$ ,  $i = 1, \dots, m$ , and is given by

$$H^*(M; \mathbb{Z}) \cong \mathbb{Z}[v_1, \dots, v_m]/\mathcal{I}, \quad \deg v_i = 2,$$

where  $\mathcal{I}$  is the ideal generated by elements of two kinds:

- (a)  $v_{i_1} \cdots v_{i_k}$ , where  $F_{i_1} \cap \cdots \cap F_{i_k} = \emptyset$  in  $P$ ;
- (b)  $\sum_{i=1}^m \langle \Lambda(F_i), \mathbf{x} \rangle v_i$  for any  $\mathbf{x} \in \mathbb{Z}^n$ .

The  $\mathbb{Z}_2$ -cohomology ring  $H^*(N; \mathbb{Z}_2)$  of a small cover  $N = N(P, \Lambda)$  has the same description, with generators  $v_i$  of degree 1.

## Theorem (Buchstaber-Erokhovets-Masuda-P-Park)

Let  $M = M(P, \Lambda)$  and  $M' = M(P', \Lambda')$  be quasitoric 6-manifolds, where  $P$  is a Pogorelov 3-polytope. The following conditions are equivalent:

- (a) There is a ring isomorphism  $\varphi: H^*(M; \mathbb{Z}) \xrightarrow{\cong} H^*(M'; \mathbb{Z})$ ;
- (b) There is a diffeomorphism  $M \cong M'$ ;
- (c) There is an equivalence of characteristic pairs  $(P, \Lambda) \sim (P', \Lambda')$ .

## Idea of proof (for both theorems).

We need to prove that a ring iso  $\varphi: H^*(N; \mathbb{Z}) \xrightarrow{\cong} H^*(N'; \mathbb{Z})$  implies an equivalence  $(P, \Lambda) \sim (P', \Lambda')$ .

An iso  $\varphi: H^*(N; \mathbb{Z}_2) \xrightarrow{\cong} H^*(N'; \mathbb{Z}_2)$  implies an iso

$\varphi: H^*(M; \mathbb{Z}_2) \xrightarrow{\cong} H^*(M'; \mathbb{Z}_2)$  (as every  $\mathbb{Z}_2$ -characteristic function of a 3-polytope lifts to a  $\mathbb{Z}$ -characteristic function!)

An iso  $\varphi: H^*(M; \mathbb{Z}_2) = \mathbb{Z}_2[P]/\mathcal{J}_\Lambda \xrightarrow{\cong} \mathbb{Z}_2[P']/\mathcal{J}_{\Lambda'} = H^*(M'; \mathbb{Z}_2)$  implies an iso

$$\begin{aligned} \psi: H^*(\mathcal{Z}_P; \mathbb{Z}_2) &= \text{Tor}_{\mathbb{Z}_2[v_1, \dots, v_m]/\mathcal{J}_\Lambda}(\mathbb{Z}_2[P]/\mathcal{J}_\Lambda, \mathbb{Z}_2) \\ &\xrightarrow{\cong} \text{Tor}_{\mathbb{Z}_2[v_1, \dots, v_m]/\mathcal{J}_{\Lambda'}}(\mathbb{Z}_2[P']/\mathcal{J}_{\Lambda'}, \mathbb{Z}_2) = H^*(\mathcal{Z}_{P'}; \mathbb{Z}_2) \end{aligned}$$

Pogorelov's conditions imply that  $\psi$  maps the set of canonical generators  $\{[u_i v_j] \in H^3(\mathcal{Z}_P): F_i \cap F_j = \emptyset\}$  bijectively to the corresponding set for  $\mathcal{Z}_{P'}$ .

This implies that  $\varphi$  maps the set  $\{[v_i] \in H^2(M)\}$  bijectively to the corresponding set for  $M'$ , giving an equivalence  $(P, \Lambda) \sim (P', \Lambda')$ . □

## Problem

*Let  $M = M(P, \Lambda)$  and  $M' = M(P', \Lambda')$  be quasitoric manifolds with isomorphic integer cohomology rings. Are they homeomorphic?*

Our result gives a positive answer in the case of quasitoric 6-manifolds over Pogorelov polytopes.

## Proposition

*6-dimensional quasitoric manifolds  $M$  and  $M'$  are diffeomorphic if there is an isomorphism  $\varphi: H^*(M; \mathbb{Z}) \xrightarrow{\cong} H^*(M'; \mathbb{Z})$  preserving the first Pontryagin class, i. e.  $\varphi(p_1(M)) = p_1(M')$ .*

We have  $p_1(M) = v_1^2 + \dots + v_m^2 \in H^4(M)$ . However, we were not able to establish the invariance of  $p_1(M)$  directly...

- [1] V. Buchstaber and T. Panov. *Toric Topology*. Mathematical Surveys and Monographs, vol. 204, American Mathematical Society, Providence, RI, 2015, 516 pages.
- [2] V. Buchstaber and T. Panov. *On manifolds defined by 4-colourings of simple 3-polytopes*. Russian Math. Surveys 71 (2016), no. 6, 1137–1139.
- [3] V. Buchstaber, N. Erokhovets, M. Masuda, T. Panov and S. Park. *Cohomological rigidity of manifolds defined by right-angled 3-dimensional polytopes*. Russian Math. Surveys 72 (2017), no. 2, 199–256.